

**Connecticut Resident Age-Adjusted Rates for 62 Cause of Death Classifications by Sex
White Residents, 1991-1992**

	Page
Connecticut Resident Age-Adjusted Mortality Rate (AAMR) Tables by Cause of Death and Sex, White Residents, 1991-1992 Rates calculated using the 1940, 1970 and 2000 standard populations	
1940 standard population	2
1970 standard population	4
2000 standard population	6
Connecticut Resident Age-Adjusted Years of Potential Life Lost Rate (YPLL) Tables by Cause of Death and Sex, White Residents, 1991-1992 Rates calculated using the 1940, 1970 and 2000 standard populations	
1940 standard population	8
1970 standard population	14
2000 standard population	20

CONNECTICUT White Resident DEATHS, 1991-1992
Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (*)
for selected causes of death by sex of decedent
*** Age-adjusted using 1940 U.S. standard population ***

	Both Sexes				Male				Female			
	*****1991-1992*****				*****1991-1992*****				*****1991-1992*****			
	No. of deaths	Crude rate	AAMR	Significance	No. of deaths	Crude rate	AAMR	Significance	No. of deaths	Crude rate	AAMR	Significance
*****Causes of Death*****												
All causes	51,930	886.4	422.5	1	435.2	25,542	898.5	551.1	0	564.9	26,388	875.0
Infectious diseases and parasitic disease (1-139)	1,332	22.7	14.1	1	12.2	772	27.2	20.3	1	17.4	560	18.6
..Septicemia (038)	692	11.8	5.0	0	5.3	293	10.3	5.8	0	6.3	399	13.2
....Septicemia-related (038)	2,521	43.0	19.4	0	20.0	1,161	40.8	23.9	0	24.2	1,360	45.1
....HIV infection (042-044)	482	8.2	7.5	1	4.9	394	13.9	12.3	1	8.4	88	2.9
All cancer (140-208)	13,173	224.8	123.6	0	124.8	6,713	236.1	148.3	0	148.9	6,460	214.2
..Oral cavity or pharynx (140.0-149.9)	229	3.9	2.4	0	2.4	140	4.9	3.6	0	3.8	89	3.0
..Colorectal cancer (153.0-154.3,154.8,159.0)	1,551	26.5	12.9	0	13.8	775	27.3	16.3	0	17.0	776	25.7
....Colon cancer (153)	1,278	21.8	10.6	0	11.6	635	22.3	13.3	0	14.0	643	21.3
....Rectal cancer (154)	248	4.2	2.1	0	2.0	128	4.5	2.7	0	2.6	120	4.0
....Pancreatic cancer (157)	689	11.8	6.0	0	6.1	294	10.3	6.5	0	7.9	395	13.1
..Lung and other respiratory cancer (160-165)	3,564	60.8	35.7	0	35.2	2,125	74.7	48.2	0	48.5	1,439	47.7
....Lung cancer (162.2-.9)	3,441	58.7	34.4	0	34.3	2,032	71.5	46.0	0	47.0	1,409	46.7
..Malignant melanoma of skin (172)	183	3.1	2.1	0	2.0	107	3.8	2.7	0	2.3	76	2.5
..Female breast cancer (174)	1,114	1,114	36.9
..Cervical cancer (180)	79	79	2.6
..Endometrial cancer (182)	76	76	2.5
..Ovarian cancer (183)	349	349	11.6
..Prostate cancer (185)	849	849	29.9	15.2	0	14.3	.	.
..Bladder cancer (188)	326	5.6	2.4	0	2.4	204	7.2	3.9	0	4.3	122	4.0
..Brain & other CNS cancer (191-192)	305	5.2	3.7	0	3.4	174	6.1	4.8	0	4.3	131	4.3
..Leukemia (204-208)	461	7.9	4.5	0	4.6	243	8.5	5.7	0	5.7	218	7.2
Diabetes mellitus (250)	882	15.1	7.3	0	7.5	394	13.9	8.6	0	8.5	488	16.2
..Diabetes-related (250)	4,215	71.9	32.4	0	34.2	1,913	67.3	39.1	0	40.8	2,302	76.3
..Diabetic ketoacidosis (DKA) (250.1)	11	0.2	.	.	.	8	0.3	.	.	.	3	0.1
Alzheimer's diseases (331.0)	371	6.3	2.0	0	2.2	140	4.9	2.3	0	2.5	231	7.7
Cardiovascular disease (390-459)	22,741	388.2	155.0	1	162.2	10,538	370.7	208.0	1	217.9	12,203	404.6
..Diseases of the heart (390-398,402,404-429)	18,103	309.0	126.3	1	132.2	8,667	304.9	173.9	1	183.6	9,436	312.9
....Heart disease-related (390-398,402,404-429)	29,725	507.4	212.6	1	219.4	14,237	500.8	285.0	0	291.9	15,488	513.6
....Coronary heart disease (CHD) (402,410-414,429.2)	13,443	229.5	93.8	1	98.8	6,623	233.0	133.5	0	140.1	6,820	226.1
....Congestive heart failure (CHF) (428.0)	841	14.4	4.3	0	4.1	298	10.5	4.8	0	5.3	543	18.0
..Hypertension with or without renal disease (401,403)	226	3.9	1.6	0	1.5	99	3.5	1.9	0	1.6	127	4.2
....Hypertension-related (401,403)	2,958	50.5	20.4	0	21.9	1,150	40.5	23.5	0	24.2	1,888	60.0
..Cerebrovascular disease (430-438)	3,247	55.4	19.6	0	20.8	1,231	43.3	22.0	0	22.2	2,016	66.8
..Atherosclerosis (440)	402	6.9	2.0	0	2.0	136	4.8	2.2	0	2.3	266	8.8
....Atherosclerosis-related (440)	1,945	33.2	10.4	1	11.7	770	27.1	13.1	0	14.1	1,175	39.0
..Aortic aneurysm (441)	407	6.9	3.0	0	2.9	237	8.3	4.7	0	4.8	170	5.6
Pneumonia & influenza (480-487)	2,140	36.5	11.5	0	12.4	938	33.0	15.6	0	16.2	1,202	39.9

(CONTINUED)

* - Crude and age-adjusted mortality rates are calculated per 100,000 population.

. - The AAMR is not reported due to a small number of deaths (<15).

1 - The 1991-1992 AAMR is significantly different (prob<0.05) from the 1989-1990 AAMR.

0 - The 1991-1992 AAMR is not significantly different (prob>0.05) from the 1989-1990 AAMR.

CONNECTICUT White Resident DEATHS, 1991-1992
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1940 U.S. standard population ***

	Both Sexes						Male						Female					
	*****1991-1992*****			1989-			*****1991-1992*****			1989-			*****1991-1992*****			1989-		
	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance
*****Causes of Death*****																		
Chronic obstructive pulmonary disease (COPD) (490-496)	2,074	35.4	14.8	0	14.9	1,068	37.6	19.6	0	19.7	1,006	33.4	11.9	0	12.1			
..COPD-related (490-496)	4,738	80.9	34.0	0	34.8	2,529	89.0	46.7	0	48.8	2,209	73.2	26.1	0	26.5			
..Asthma (493)	74	1.3	0.7	0	0.8	21	0.7	0.6	0	0.8	53	1.8	0.7	0	0.9			
Chronic liver disease and cirrhosis (571)	540	9.2	6.6	0	7.2	340	12.0	9.4	0	10.0	200	6.6	4.2	0	4.8			
Nephritis, nephrotic syndrome/nephrosis (580-589)	566	9.7	3.9	0	3.5	292	10.3	5.4	0	4.7	274	9.1	3.0	0	2.8			
..Nephritis-related (580-589)	2,901	49.5	21.6	0	23.4	1,525	53.6	29.4	0	32.2	1,376	45.6	16.4	0	17.5			
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	632	10.8	4.4	0	3.8	323	11.4	6.0	0	4.9	309	10.2	3.4	0	3.1			
Unintentional injuries (E800-E949)	1,551	26.5	20.4	1	22.8	990	34.8	30.3	0	34.4	561	18.6	10.9	0	11.6			
..Motor vehicle accidents (E810-E825)	581	9.9	9.8	1	12.7	393	13.8	14.1	1	19.2	188	6.2	5.5	0	6.3			
..Falls and fall related injuries (E880-E888)	340	5.8	2.3	0	2.3	163	5.7	3.3	0	3.1	177	5.9	1.5	0	1.7			
..Drowning (E830,E832,E910)	67	1.1	1.0	0	1.3	55	1.9	1.8	0	2.2	121	0.4	.1	0	0.5			
..Residential fires (E890-E899)	53	0.9	0.8	0	1.0	26	0.9	0.8	0	1.3	27	0.9	0.8	0	0.6			
Suicide (E950-E959)	611	10.4	9.5	0	8.2	501	17.6	16.1	0	13.6	110	3.6	3.3	0	3.0			
Homicide and legal intervention (E960-E978)	204	3.5	3.7	0	3.5	155	5.5	5.7	0	5.0	49	1.6	1.7	0	2.0			
..Homicide (E960-E969)	203	3.5	3.7	0	3.5	154	5.4	5.7	0	5.0	49	1.6	1.7	0	2.0			
All injury (E800-E999)	2,443	41.7	34.8	0	35.0	1,696	59.7	53.7	0	54.0	747	24.8	16.7	0	16.8			
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	569	9.7	9.6	1	12.6	383	13.5	13.7	1	18.9	186	6.2	5.4	0	6.3			
..All falls (E880-E886,E888,E957,E968.1,E987)	283	4.8	2.2	0	2.1	146	5.1	3.1	0	2.8	137	4.5	1.4	0	1.3			
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	419	7.2	7.1	0	6.2	369	13.0	12.7	0	10.4	50	1.7	1.7	0	2.1			
....Suicide (E955.0-.4)	267	4.6	4.2	0	3.5	241	8.5	7.8	0	6.2	26	0.9	0.8	0	1.0			
....Homicide (E965.0-.4)	136	2.3	2.6	0	2.2	117	4.1	4.5	0	3.3	19	0.6	0.7	0	1.0			
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	360	6.1	5.6	1	2.8	267	9.4	8.5	1	4.2	93	3.1	2.7	1	1.5			
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	314	5.4	4.3	0	4.5	227	8.0	6.6	0	6.6	87	2.9	2.2	0	2.6			
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	239	4.1	3.8	1	1.1	175	6.2	5.7	1	1.5	64	2.1	2.0	1	0.7			

* - Crude and age-adjusted mortality rates are calculated per 100,000 population.

. - The AAMR is not reported due to a small number of deaths (<15).

1 - The 1991-1992 AAMR is significantly different (prob<0.05) from the 1989-1990 AAMR.

0 - The 1991-1992 AAMR is not significantly different (prob>0.05) from the 1989-1990 AAMR.

CONNECTICUT White Resident DEATHS, 1991-1992
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1970 U.S. standard population ***

	Both Sexes						Male						Female					
	*****1991-1992*****			1989-			*****1991-1992*****			1989-			*****1991-1992*****			1989-		
	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance	No. of deaths	Crude rate	Signif-icance
*****Causes of Death*****																		
Chronic obstructive pulmonary disease (COPD) (490-496)	2,074	35.4	22.8	0	22.3	1,068	37.6	31.4	0	30.9	1,006	33.4	18.0	0	17.6			
..COPD-related (490-496)	4,738	80.9	52.2	0	52.6	2,529	89.0	74.6	0	77.3	2,209	73.2	39.4	0	38.8			
..Asthma (493)	74	1.3	0.9	0	1.1	21	0.7	0.7	0	1.0	53	1.8	1.0	0	1.1			
Chronic liver disease and cirrhosis (571)	540	9.2	7.4	0	8.2	340	12.0	10.5	0	11.4	200	6.6	4.8	0	5.6			
Nephritis, nephrotic syndrome/nephrosis (580-589)	566	9.7	6.1	0	5.5	292	10.3	8.7	0	7.6	274	9.1	4.6	0	4.3			
..Nephritis-related (580-589)	2,901	49.5	32.2	0	34.6	1,525	53.6	45.3	0	49.2	1,376	45.6	24.5	0	25.9			
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	632	10.8	6.8	0	5.9	323	11.4	9.6	0	8.0	309	10.2	5.2	0	4.8			
Unintentional injuries (E800-E949)	1,551	26.5	21.8	1	24.3	990	34.8	31.9	0	35.8	561	18.6	12.7	0	13.6			
..Motor vehicle accidents (E810-E825)	581	9.9	9.5	1	12.4	393	13.8	13.6	1	18.3	188	6.2	5.5	0	6.4			
..Falls and fall related injuries (E880-E888)	340	5.8	3.6	0	3.6	163	5.7	4.9	0	4.4	177	5.9	2.6	0	2.9			
..Drowning (E830,E832,E910)	67	1.1	1.0	0	1.3	55	1.9	1.8	0	2.2	121	0.4	.1	0	0.5			
..Residential fires (E890-E899)	53	0.9	0.9	0	1.0	26	0.9	0.9	0	1.4	27	0.9	0.9	0	0.7			
Suicide (E950-E959)	611	10.4	9.1	0	7.9	501	17.6	15.6	0	13.3	110	3.6	3.1	0	2.9			
Homicide and legal intervention (E960-E978)	204	3.5	3.4	0	3.2	155	5.5	5.2	0	4.6	49	1.6	1.6	0	1.9			
..Homicide (E960-E969)	203	3.5	3.4	0	3.2	154	5.4	5.2	0	4.6	49	1.6	1.6	0	1.8			
All injury (E800-E999)	2,443	41.7	35.4	0	36.0	1,696	59.7	54.2	0	54.6	747	24.8	18.0	0	18.6			
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	569	9.7	9.3	1	12.2	383	13.5	13.2	1	18.0	186	6.2	5.5	0	6.4			
..All falls (E880-E886,E888,E957,E968.1,E987)	283	4.8	3.1	0	3.0	146	5.1	4.4	0	3.8	137	4.5	2.2	0	2.3			
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	419	7.2	6.7	0	5.8	369	13.0	12.2	0	10.0	50	1.7	1.5	0	1.9			
....Suicide (E955.0-.4)	267	4.6	4.0	0	3.4	241	8.5	7.7	0	6.1	26	0.9	0.7	0	0.9			
....Homicide (E965.0-.4)	136	2.3	2.4	0	2.0	117	4.1	4.1	0	3.0	19	0.6	0.6	0	1.0			
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	360	6.1	5.0	1	2.7	267	9.4	7.5	1	4.0	93	3.1	2.6	1	1.5			
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	314	5.4	4.4	0	4.7	227	8.0	6.8	0	7.0	87	2.9	2.3	0	2.7			
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	239	4.1	3.3	1	1.0	175	6.2	4.8	1	1.3	64	2.1	1.8	1	0.7			

* - Crude and age-adjusted mortality rates are calculated per 100,000 population.

. - The AAMR is not reported due to a small number of deaths (<15).

1 - The 1991-1992 AAMR is significantly different (prob<0.05) from the 1989-1990 AAMR.

0 - The 1991-1992 AAMR is not significantly different (prob>0.05) from the 1989-1990 AAMR.

CONNECTICUT White Resident DEATHS, 1991-1992
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 2000 U.S. standard population ***

	Both Sexes				Male				Female				1989- 1990	
	*****1991-1992*****		1989-		*****1991-1992*****		1989-		*****1991-1992*****		1989-			
	No. of deaths	Crude rate	Signif-icance	AAMR	No. of deaths	Crude rate	Signif-icance	AAMR	No. of deaths	Crude rate	Signif-icance	AAMR		
*****Causes of Death*****														
Chronic obstructive pulmonary disease (COPD) (490-496)	2,074	35.4	31.8	0	30.2	1,068	37.6	45.6	0	43.8	1,006	33.4	24.7	
..COPD-related (490-496)	4,738	80.9	72.9	0	72.0	2,529	89.0	108.0	0	110.8	2,209	73.2	54.5	
..Asthma (493)	74	1.3	1.2	0	1.3	21	0.7	0.8	0	1.3	53	1.8	1.3	
Chronic liver disease and cirrhosis (571)	540	9.2	8.7	0	9.6	340	12.0	12.4	0	13.4	200	6.6	5.6	
Nephritis, nephrotic syndrome/nephrosis (580-589)	566	9.7	8.9	0	8.0	292	10.3	13.0	0	11.4	274	9.1	6.7	
..Nephritis-related (580-589)	2,901	49.5	45.2	0	48.4	1,525	53.6	65.5	0	70.6	1,376	45.6	34.2	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	632	10.8	9.9	0	8.6	323	11.4	14.4	0	12.0	309	10.2	7.6	
Unintentional injuries (E800-E949)	1,551	26.5	25.3	0	27.4	990	34.8	36.7	0	40.0	561	18.6	15.5	
..Motor vehicle accidents (E810-E825)	581	9.9	9.6	1	12.2	393	13.8	13.7	1	18.1	188	6.2	5.8	
..Falls and fall related injuries (E880-E888)	340	5.8	5.4	0	5.4	163	5.7	7.3	0	6.2	177	5.9	4.2	
..Drowning (E830,E832,E910)	67	1.1	1.1	0	1.4	55	1.9	1.9	0	2.3	121	0.4	.	
..Residential fires (E890-E899)	53	0.9	0.9	0	1.1	26	0.9	1.0	0	1.5	27	0.9	0.9	
Suicide (E950-E959)	611	10.4	10.0	0	8.7	501	17.6	17.3	0	15.0	110	3.6	3.5	
Homicide and legal intervention (E960-E978)	204	3.5	3.4	0	3.3	155	5.5	5.2	0	4.6	49	1.6	1.6	
..Homicide (E960-E969)	203	3.5	3.4	0	3.3	154	5.4	5.1	0	4.6	49	1.6	1.6	
All injury (E800-E999)	2,443	41.7	39.9	0	40.1	1,696	59.7	60.9	0	60.6	747	24.8	21.4	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	569	9.7	9.4	1	12.0	383	13.5	13.3	1	17.8	186	6.2	5.7	
..All falls (E880-E886,E888,E957,E968.1,E987)	283	4.8	4.5	0	4.4	146	5.1	6.3	0	5.1	137	4.5	3.3	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	419	7.2	6.9	0	6.2	369	13.0	12.6	0	10.7	50	1.7	1.6	
....Suicide (E955.0-.4)	267	4.6	4.4	0	3.7	241	8.5	8.3	0	6.9	26	0.9	0.8	
....Homicide (E965.0-.4)	136	2.3	2.3	0	2.0	117	4.1	3.9	0	2.9	19	0.6	0.6	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	360	6.1	5.9	1	3.1	267	9.4	9.0	1	4.7	93	3.1	3.0	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-3,790.3,E860)	314	5.4	5.2	0	5.4	227	8.0	8.1	0	8.0	87	2.9	2.6	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-5,E962.0,E980.0-.5)	239	4.1	3.9	1	1.2	175	6.2	5.7	1	1.7	64	2.1	2.1	

* - Crude and age-adjusted mortality rates are calculated per 100,000 population.

. - The AAMR is not reported due to a small number of deaths (<15).

1 - The 1991-1992 AAMR is significantly different (prob<0.05) from the 1989-1990 AAMR.

0 - The 1991-1992 AAMR is not significantly different (prob>0.05) from the 1989-1990 AAMR.

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1940 U.S. standard population ***

	Both Sexes											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	74	19	42	398	705	8.3	10.3	0	12.5	15.6	0	
Chronic liver disease and cirrhosis (571)	540	291	435	4,258	7,888	72.9	71.4	0	124.2	127.8	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	566	70	190	822	2,022	14.9	17.2	0	30.5	32.5	0	
..Nephritis-related (580-589)	2,901	424	1,115	5,253	12,576	95.4	104.4	0	191.0	210.5	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	632	77	212	850	2,207	15.3	17.2	0	33.0	33.4	0	
Unintentional injuries (E800-E949)	1,551	904	1,091	27,756	37,684	588.4	706.5	1	738.0	873.0	1	
..Motor vehicle accidents (E810-E825)	581	463	513	15,568	20,443	344.9	472.2	1	421.9	571.8	1	
..Falls and fall related injuries (E880-E888)	340	54	100	880	1,635	16.9	26.8	0	27.3	37.0	0	
..Drowning (E830,E832,E910)	67	49	59	1,577	2,112	33.4	49.4	0	41.4	60.1	0	
..Residential fires (E890-E899)	53	35	47	1,237	1,637	27.3	30.9	0	33.3	38.1	0	
Suicide (E950-E959)	611	498	553	13,765	19,038	273.4	223.6	0	351.6	290.7	0	
Homicide and legal intervention (E960-E978)	204	191	197	6,636	8,566	143.2	126.2	0	173.4	154.3	0	
..Homicide (E960-E969)	203	190	196	6,598	8,518	142.4	125.7	0	172.6	153.7	0	
All injury (E800-E999)	2,443	1,665	1,914	50,224	68,076	1042.4	1074.4	0	1310.8	1341.2	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	569	454	503	15,345	20,123	339.6	468.1	1	415.1	566.5	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	283	64	106	1,200	2,040	22.7	26.8	0	34.3	37.0	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	419	360	391	11,345	15,103	240.2	200.6	0	298.3	251.2	0	
....Suicide (E955.0-.4)	267	215	243	6,198	8,493	125.8	97.8	0	160.4	126.5	0	
....Homicide (E965.0-.4)	136	132	134	4,660	5,990	103.0	81.6	0	124.3	99.2	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	360	323	339	8,848	12,158	162.5	68.5	1	209.9	91.5	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	314	220	285	3,920	6,458	66.4	58.3	0	102.5	96.9	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	239	228	232	6,495	8,790	118.1	25.3	1	150.7	33.8	1	

(CONTINUED)

* - Age-adjusted YPLL are calculated per 100,000 population.
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1940 U.S. standard population ***

	Male											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	21	11	17	198	343	8.0	11.2	0	12.3	16.2	0	
Chronic liver disease and cirrhosis (571)	340	206	286	3,015	5,460	105.3	101.6	0	178.3	180.3	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	292	44	109	490	1,203	18.0	22.1	0	37.9	39.6	0	
..Nephritis-related (580-589)	1,525	264	647	3,208	7,516	115.0	129.9	0	235.2	261.8	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	323	47	121	498	1,303	18.2	21.8	0	40.6	40.1	0	
Unintentional injuries (E800-E949)	990	687	800	21,359	28,781	893.0	1086.1	1	1119.0	1342.6	1	
..Motor vehicle accidents (E810-E825)	393	339	363	11,674	15,189	510.8	716.9	1	622.4	868.3	1	
..Falls and fall related injuries (E880-E888)	163	41	72	728	1,285	28.3	44.8	0	44.3	60.2	0	
..Drowning (E830,E832,E910)	55	43	51	1,385	1,850	57.6	77.7	0	71.6	95.7	0	
..Residential fires (E890-E899)	26	17	22	621	818	27.0	41.8	0	33.1	51.2	0	
Suicide (E950-E959)	501	405	455	11,383	15,693	453.5	372.8	0	582.9	482.7	0	
Homicide and legal intervention (E960-E978)	155	147	152	5,045	6,532	216.6	185.0	0	263.3	225.2	0	
..Homicide (E960-E969)	154	146	151	5,007	6,485	215.1	185.0	0	261.6	225.2	0	
All injury (E800-E999)	1,696	1,287	1,455	39,198	52,898	1615.0	1674.9	0	2030.9	2090.1	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	383	331	354	11,464	14,891	500.9	708.7	1	609.7	857.9	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	146	47	75	923	1,528	35.4	47.4	0	52.7	63.3	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	369	314	343	9,955	13,243	421.2	330.2	0	523.5	414.2	0	
....Suicide (E955.0-.4)	241	191	218	5,558	7,665	226.8	165.7	0	289.2	215.2	0	
....Homicide (E965.0-.4)	117	113	115	3,988	5,128	175.2	126.3	0	211.6	153.1	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	267	247	257	6,933	9,463	252.1	107.1	1	324.5	141.0	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	227	165	204	2,933	4,790	100.6	83.7	0	155.4	139.3	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	175	171	172	5,063	6,775	182.3	36.9	1	230.5	48.6	1	

(CONTINUED)

* - Age-adjusted YPLL are calculated per 100,000 population.
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1940 U.S. standard population ***

	Female											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	53	8	25	200	363	8.7	9.2	0	12.8	14.8	0	
Chronic liver disease and cirrhosis (571)	200	85	149	1,243	2,428	41.5	42.3	0	72.4	77.8	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	274	26	81	332	820	11.9	12.2	0	23.7	25.4	0	
..Nephritis-related (580-589)	1,376	160	468	2,045	5,060	76.7	79.6	0	150.4	162.7	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	309	30	91	352	905	12.6	12.6	0	26.0	26.7	0	
Unintentional injuries (E800-E949)	561	217	291	6,398	8,963	276.8	319.3	0	350.7	396.1	0	
..Motor vehicle accidents (E810-E825)	188	124	150	3,894	5,254	173.8	221.5	0	215.7	269.0	0	
..Falls and fall related injuries (E880-E888)	177	13	28	153	350	5.5	8.3	0	10.6	13.6	0	
..Drowning (E830,E832,E910)	12	6	8	192	262	.	20.9	.	.	24.2	.	
..Residential fires (E890-E899)	27	18	25	616	819	27.5	19.6	0	33.5	24.5	0	
Suicide (E950-E959)	110	93	98	2,383	3,345	90.5	73.1	0	118.4	98.3	0	
Homicide and legal intervention (E960-E978)	49	44	45	1,591	2,034	67.7	65.7	0	81.4	81.6	0	
..Homicide (E960-E969)	49	44	45	1,591	2,034	67.7	64.8	0	81.4	80.5	0	
All injury (E800-E999)	747	378	459	11,026	15,179	457.6	463.3	0	579.8	583.2	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	186	123	149	3,882	5,232	173.3	221.5	0	214.9	268.7	0	
..All falls (E880-E886,E888,E957,E968.1,E987)	137	17	31	278	513	9.8	5.8	0	16.0	10.4	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	50	46	48	1,390	1,860	54.5	68.8	0	68.4	86.5	0	
....Suicide (E955.0-.4)	26	24	25	640	888	22.9	29.3	0	29.9	37.8	0	
....Homicide (E965.0-.4)	19	19	19	673	863	28.6	35.8	0	34.5	44.1	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	93	76	82	1,915	2,695	72.8	29.7	1	95.5	42.0	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	87	55	81	988	1,668	33.0	33.6	0	51.3	56.3	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	64	57	60	1,433	2,015	53.6	13.7	1	70.5	19.2	1	

* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1970 U.S. standard population ***

	Both Sexes											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	74	19	42	398	705	8.0	10.3	0	12.4	15.8	0	
Chronic liver disease and cirrhosis (571)	540	291	435	4,258	7,888	67.5	67.5	0	120.4	126.4	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	566	70	190	822	2,022	14.4	17.3	0	32.0	34.6	0	
..Nephritis-related (580-589)	2,901	424	1,115	5,253	12,576	93.0	102.3	0	200.3	221.0	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	632	77	212	850	2,207	14.9	17.5	0	34.9	35.8	0	
Unintentional injuries (E800-E949)	1,551	904	1,091	27,756	37,684	558.6	679.8	1	694.2	830.1	1	
..Motor vehicle accidents (E810-E825)	581	463	513	15,568	20,443	328.0	452.6	1	396.2	540.1	1	
..Falls and fall related injuries (E880-E888)	340	54	100	880	1,635	15.9	27.3	0	26.8	38.0	0	
..Drowning (E830,E832,E910)	67	49	59	1,577	2,112	32.4	49.3	0	39.6	58.9	0	
..Residential fires (E890-E899)	53	35	47	1,237	1,637	28.8	31.9	0	34.7	38.8	0	
..Suicide (E950-E959)	611	498	553	13,765	19,038	247.7	203.2	0	316.9	263.2	0	
Homicide and legal intervention (E960-E978)	204	191	197	6,636	8,566	133.8	119.2	0	159.4	143.4	0	
..Homicide (E960-E969)	203	190	196	6,598	8,518	133.2	118.8	0	158.7	143.0	0	
All injury (E800-E999)	2,443	1,665	1,914	50,224	68,076	973.0	1018.2	0	1211.7	1257.1	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	569	454	503	15,345	20,123	322.9	448.7	1	389.4	535.0	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	283	64	106	1,200	2,040	21.1	27.0	0	33.0	37.6	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	419	360	391	11,345	15,103	222.1	186.2	0	272.6	230.6	0	
...Suicide (E955.0-.4)	267	215	243	6,198	8,493	114.3	89.5	0	144.8	115.2	0	
...Homicide (E965.0-.4)	136	132	134	4,660	5,990	97.1	76.4	0	115.1	91.4	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	360	323	339	8,848	12,158	141.5	60.7	1	181.5	81.2	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	314	220	285	3,920	6,458	59.3	54.4	0	94.4	93.3	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	239	228	232	6,495	8,790	101.4	20.9	1	128.1	28.1	1	

(CONTINUED)

* - Age-adjusted YPLL are calculated per 100,000 population.
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1970 U.S. standard population ***

	Male											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	21	11	17	198	343	7.6	11.0	0	12.1	16.0	0	
Chronic liver disease and cirrhosis (571)	340	206	286	3,015	5,460	98.7	95.7	0	173.5	177.8	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	292	44	109	490	1,203	17.2	22.6	0	39.7	42.2	0	
..Nephritis-related (580-589)	1,525	264	647	3,208	7,516	111.1	126.8	0	245.9	274.3	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	323	47	121	498	1,303	17.5	22.5	0	42.9	43.1	0	
Unintentional injuries (E800-E949)	990	687	800	21,359	28,781	843.1	1037.1	1	1046.4	1266.5	1	
..Motor vehicle accidents (E810-E825)	393	339	363	11,674	15,189	485.9	682.5	1	584.1	814.3	1	
..Falls and fall related injuries (E880-E888)	163	41	72	728	1,285	26.4	45.6	0	43.2	61.5	0	
..Drowning (E830,E832,E910)	55	43	51	1,385	1,850	55.7	76.5	0	68.2	92.6	0	
..Residential fires (E890-E899)	26	17	22	621	818	28.8	43.7	0	34.7	52.9	0	
..Suicide (E950-E959)	501	405	455	11,383	15,693	412.2	339.1	0	527.3	437.3	0	
Homicide and legal intervention (E960-E978)	155	147	152	5,045	6,532	201.3	174.9	0	241.1	209.7	0	
..Homicide (E960-E969)	154	146	151	5,007	6,485	200.2	174.9	0	239.7	209.7	0	
All injury (E800-E999)	1,696	1,287	1,455	39,198	52,898	1503.3	1578.6	0	1872.7	1948.2	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	383	331	354	11,464	14,891	476.2	674.8	1	571.5	804.4	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	146	47	75	923	1,528	33.2	47.6	0	51.0	63.8	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	369	314	343	9,955	13,243	390.1	307.6	0	479.7	381.8	0	
...Suicide (E955.0-.4)	241	191	218	5,558	7,665	207.0	152.2	0	262.7	196.9	0	
...Homicide (E965.0-.4)	117	113	115	3,988	5,128	164.6	118.4	0	195.6	141.2	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	267	247	257	6,933	9,463	218.1	94.6	1	278.4	124.5	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	227	165	204	2,933	4,790	90.4	77.6	0	143.7	134.0	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	175	171	172	5,063	6,775	155.2	29.7	1	193.7	39.2	1	

(CONTINUED)

* - Age-adjusted YPLL are calculated per 100,000 population.
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 1970 U.S. standard population ***

	Female											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	53	8	25	200	363	8.4	9.5	0	12.7	15.4	0	
Chronic liver disease and cirrhosis (571)	200	85	149	1,243	2,428	37.5	40.2	0	69.8	77.7	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	274	26	81	332	820	11.7	12.0	0	24.9	27.0	0	
..Nephritis-related (580-589)	1,376	160	468	2,045	5,060	76.0	78.5	0	158.9	172.0	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	309	30	91	352	905	12.5	12.4	0	27.7	28.6	0	
Unintentional injuries (E800-E949)	561	217	291	6,398	8,963	267.0	314.6	0	335.8	386.5	0	
..Motor vehicle accidents (E810-E825)	188	124	150	3,894	5,254	164.9	216.7	0	202.6	259.6	0	
..Falls and fall related injuries (E880-E888)	177	13	28	153	350	5.3	8.4	0	10.9	14.2	0	
..Drowning (E830,E832,E910)	12	6	8	192	262	.	21.7	.	.	24.8	.	
..Residential fires (E890-E899)	27	18	25	616	819	28.8	19.7	0	34.6	24.4	0	
Suicide (E950-E959)	110	93	98	2,383	3,345	80.4	65.9	0	104.8	88.9	0	
Homicide and legal intervention (E960-E978)	49	44	45	1,591	2,034	64.1	61.7	0	75.6	75.5	0	
..Homicide (E960-E969)	49	44	45	1,591	2,034	64.1	61.0	0	75.6	74.6	0	
All injury (E800-E999)	747	378	459	11,026	15,179	430.3	446.6	0	540.4	556.8	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	186	123	149	3,882	5,232	164.4	216.7	0	201.8	259.2	0	
..All falls (E880-E886,E888,E957,E968.1,E987)	137	17	31	278	513	8.8	6.0	0	15.2	11.2	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	50	46	48	1,390	1,860	49.5	62.6	0	61.3	77.7	0	
....Suicide (E955.0-.4)	26	24	25	640	888	19.6	26.1	0	25.5	33.4	0	
....Homicide (E965.0-.4)	19	19	19	673	863	27.2	33.3	0	32.2	40.4	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	93	76	82	1,915	2,695	64.8	26.6	1	84.7	38.0	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	87	55	81	988	1,668	29.0	31.8	0	47.0	54.7	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	64	57	60	1,433	2,015	47.4	12.1	1	62.2	17.2	1	

* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 2000 U.S. standard population ***

	Both Sexes											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	74	19	42	398	705	8.0	10.7	0	12.6	16.6	0	
Chronic liver disease and cirrhosis (571)	540	291	435	4,258	7,888	85.4	84.5	0	144.3	149.4	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	566	70	190	822	2,022	16.4	17.9	0	35.2	35.7	0	
..Nephritis-related (580-589)	2,901	424	1,115	5,253	12,576	104.9	116.2	0	218.7	242.4	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	632	77	212	850	2,207	17.0	18.1	0	38.1	37.0	0	
Unintentional injuries (E800-E949)	1,551	904	1,091	27,756	37,684	541.5	639.1	1	679.3	787.8	1	
..Motor vehicle accidents (E810-E825)	581	463	513	15,568	20,443	305.2	414.8	1	371.1	497.9	1	
..Falls and fall related injuries (E880-E888)	340	54	100	880	1,635	17.1	25.5	0	28.7	36.3	0	
..Drowning (E830,E832,E910)	67	49	59	1,577	2,112	31.5	46.6	0	38.9	56.3	0	
..Residential fires (E890-E899)	53	35	47	1,237	1,637	25.5	28.7	0	31.2	35.4	0	
Suicide (E950-E959)	611	498	553	13,765	19,038	260.4	214.5	0	334.9	279.4	0	
Homicide and legal intervention (E960-E978)	204	191	197	6,636	8,566	127.1	115.8	0	152.3	140.5	0	
..Homicide (E960-E969)	203	190	196	6,598	8,518	126.5	115.2	0	151.6	139.8	0	
All injury (E800-E999)	2,443	1,665	1,914	50,224	68,076	967.0	987.8	0	1214.5	1231.1	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	569	454	503	15,345	20,123	300.6	411.2	1	365.0	493.1	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	283	64	106	1,200	2,040	22.9	25.4	0	35.6	36.1	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	419	360	391	11,345	15,103	217.1	185.9	0	268.3	232.2	0	
....Suicide (E955.0-.4)	267	215	243	6,198	8,493	116.8	94.1	0	149.0	121.9	0	
....Homicide (E965.0-.4)	136	132	134	4,660	5,990	90.8	73.0	0	108.2	87.8	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	360	323	339	8,848	12,158	163.3	68.7	1	209.7	91.8	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	314	220	285	3,920	6,458	76.0	68.6	0	116.3	112.4	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	239	228	232	6,495	8,790	118.1	26.4	1	149.3	35.1	1	

(CONTINUED)

* - Age-adjusted YPLL are calculated per 100,000 population.
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT White Resident DEATHS, 1991-1992
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (*)
 for selected causes of death by sex of decedent
 *** Age-adjusted using 2000 U.S. standard population ***

	Male											
	Number of Deaths, 1991-1992			Years of life lost, 1991-1992		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1991-1992	1989-1990	Significance	1991-1992	1989-1990	Significance	
*****Causes of Death*****												
..Asthma (493)	21	11	17	198	343	7.9	11.3	0	12.5	16.6	0	
Chronic liver disease and cirrhosis (571)	340	206	286	3,015	5,460	124.7	119.9	0	208.5	210.2	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	292	44	109	490	1,203	19.8	23.0	0	43.7	43.1	0	
..Nephritis-related (580-589)	1,525	264	647	3,208	7,516	128.3	146.3	0	271.9	303.9	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	323	47	121	498	1,303	20.1	23.0	0	46.9	44.3	0	
Unintentional injuries (E800-E949)	990	687	800	21,359	28,781	821.5	985.0	1	1028.9	1213.6	1	
..Motor vehicle accidents (E810-E825)	393	339	363	11,674	15,189	450.3	630.3	1	544.5	756.1	1	
..Falls and fall related injuries (E880-E888)	163	41	72	728	1,285	28.1	42.6	0	45.7	58.7	0	
..Drowning (E830,E832,E910)	55	43	51	1,385	1,850	54.7	73.7	0	67.8	90.3	0	
..Residential fires (E890-E899)	26	17	22	621	818	25.3	38.2	0	31.0	46.8	0	
Suicide (E950-E959)	501	405	455	11,383	15,693	429.7	356.1	0	552.8	462.1	0	
Homicide and legal intervention (E960-E978)	155	147	152	5,045	6,532	191.2	167.0	0	230.1	201.6	0	
..Homicide (E960-E969)	154	146	151	5,007	6,485	190.0	167.0	0	228.7	201.6	0	
All injury (E800-E999)	1,696	1,287	1,455	39,198	52,898	1495.5	1539.1	0	1878.2	1916.6	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	383	331	354	11,464	14,891	441.8	623.1	1	533.4	746.8	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	146	47	75	923	1,528	35.4	44.9	0	54.2	61.3	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	369	314	343	9,955	13,243	378.9	304.3	0	469.2	381.0	0	
...Suicide (E955.0-.4)	241	191	218	5,558	7,665	209.1	159.0	0	267.1	207.1	0	
...Homicide (E965.0-.4)	117	113	115	3,988	5,128	153.8	111.5	0	183.6	133.6	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	267	247	257	6,933	9,463	253.9	106.1	1	324.5	139.6	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	227	165	204	2,933	4,790	115.6	97.7	0	176.7	160.7	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	175	171	172	5,063	6,775	181.9	38.2	1	227.6	49.8	1	

(CONTINUED)

* - Age-adjusted YPLL are calculated per 100,000 population.
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

