

**Connecticut Resident Age-Adjusted Rates for 62 Cause of Death Classifications by Sex  
White Residents, 1996-1998**

	Page
Connecticut Resident Age-Adjusted Mortality Rate (AAMR) Tables by Cause of Death and Sex, White Residents, 1996-1998 Rates calculated using the 1940, 1970 and 2000 standard populations	
1940 standard population	2
1970 standard population	4
2000 standard population	6
Connecticut Resident Age-Adjusted Years of Potential Life Lost Rate (YPLL) Tables by Cause of Death and Sex, White Residents, 1996-1998 Rates calculated using the 1940, 1970 and 2000 standard populations	
1940 standard population	8
1970 standard population	14
2000 standard population	20











CONNECTICUT White Resident DEATHS, 1996-1998  
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Both Sexes				Male				Female			
	*****1996-1998*****				*****1996-1998*****				*****1996-1998*****			
	No. of deaths	Crude rate	Signif-icance	AAMR	No. of deaths	Crude rate	Signif-icance	AAMR	No. of deaths	Crude rate	Signif-icance	AAMR
<b>*****Causes of Death*****</b>												
Chronic obstructive pulmonary disease (COPD) (490-496)	3,600	41.6	34.3	1	29.9	1,645	39.0	42.0	0	43.2	1,955	44.0
..COPD-related (490-496)	8,237	95.1	78.6	1	71.5	4,035	95.7	102.6	0	108.8	4,202	94.6
..Asthma (493)	139	1.6	1.4	0	1.2	39	0.9	1.0	0	1.0	100	2.3
Chronic liver disease and cirrhosis (571)	820	9.5	8.6	0	9.4	509	12.1	11.9	0	13.1	311	7.0
Nephritis, nephrotic syndrome/nephrosis (580-589)	951	11.0	9.1	0	8.4	460	10.9	12.2	0	11.9	491	11.1
..Nephritis-related (580-589)	4,241	49.0	40.8	1	47.0	2,253	53.5	58.1	1	68.4	1,988	44.7
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	1,121	12.9	10.7	1	9.1	556	13.2	14.7	0	12.9	565	12.7
Unintentional injuries (E800-E949)	2,836	32.8	30.3	1	26.5	1,789	42.4	43.5	1	38.8	1,047	23.6
..Motor vehicle accidents (E810-E825)	886	10.2	10.2	0	11.4	561	13.3	13.5	1	16.5	325	7.3
..Falls and fall related injuries (E880-E888)	601	6.9	5.8	0	5.4	314	7.4	8.2	0	6.7	287	6.5
..Drowning (E830,E832,E910)	70	0.8	0.8	1	1.3	58	1.4	1.4	1	2.3	12	0.3
..Residential fires (E890-E899)	80	0.9	0.9	0	1.1	45	1.1	1.1	0	1.4	35	0.8
Suicide (E950-E959)	759	8.8	8.4	0	9.3	586	13.9	13.7	0	16.0	173	3.9
Homicide and legal intervention (E960-E978)	219	2.5	2.6	0	3.3	158	3.7	3.8	0	4.8	61	1.4
..Homicide (E960-E969)	216	2.5	2.6	1	3.3	155	3.7	3.8	0	4.8	61	1.4
All injury (E800-E999)	3,915	45.2	42.5	0	40.0	2,597	61.6	62.5	0	61.0	1,318	29.7
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	865	10.0	9.9	0	11.2	546	13.0	13.1	1	16.1	319	7.2
..All falls (E880-E886,E888,E957,E968.1,E987)	441	5.1	4.3	0	4.5	255	6.1	6.4	0	5.6	186	4.2
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	422	4.9	4.8	1	6.5	370	8.8	8.8	1	11.4	52	1.2
....Suicide (E955.0-.4)	288	3.3	3.2	1	4.0	260	6.2	6.1	0	7.5	28	0.6
....Homicide (E965.0-.4)	121	1.4	1.5	1	2.1	98	2.3	2.4	0	3.3	23	0.5
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	782	9.0	8.7	1	3.8	566	13.4	12.8	1	5.6	216	4.7
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	428	4.9	4.7	0	5.1	311	7.4	7.2	0	7.6	117	2.6
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	677	7.8	7.5	1	1.9	479	11.4	10.8	1	2.7	198	4.5

\* - Crude and age-adjusted mortality rates are calculated per 100,000 population.

. - The AAMR is not reported due to a small number of deaths (<15).

1 - The 1996-1998 AAMR is significantly different ( $\text{prob} < 0.05$ ) from the 1989-1991 AAMR.

0 - The 1996-1998 AAMR is not significantly different ( $\text{prob} \geq 0.05$ ) from the 1989-1991 AAMR.



**CONNECTICUT White Resident DEATHS, 1996-1998**  
**Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)**  
**for selected causes of death by sex of decedent**  
\*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Both Sexes											
	Number of Deaths, 1996-1998			Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1996-1998	1989-1991	Significance	1996-1998	1989-1991	Significance	
<b>*****Causes of Death*****</b>												
..Asthma (493)	139	65	82	1,148	1,865	14.7	9.0	0	21.7	13.7	0	
Chronic liver disease and cirrhosis (571)	820	421	623	6,015	11,215	65.9	70.9	0	114.7	125.5	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	951	108	267	1,511	3,254	18.9	17.3	0	34.9	33.7	0	
..Nephritis-related (580-589)	4,241	607	1,422	7,623	17,200	89.7	105.1	0	177.6	208.2	1	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	1,121	136	333	1,950	4,143	24.6	17.5	0	44.7	35.0	0	
Unintentional injuries (E800-E949)	2,836	1,607	1,889	46,371	63,701	675.0	674.7	0	854.1	834.9	0	
..Motor vehicle accidents (E810-E825)	886	678	759	21,357	28,505	339.2	439.1	1	418.5	531.9	1	
..Falls and fall related injuries (E880-E888)	601	87	165	1,865	3,165	25.4	22.2	0	37.2	32.1	0	
..Drowning (E830,E832,E910)	70	59	63	2,168	2,778	33.7	44.8	0	40.3	54.6	0	
..Residential fires (E890-E899)	80	52	64	1,510	2,080	21.6	34.8	0	27.3	42.4	0	
Suicide (E950-E959)	759	617	681	15,768	22,273	213.9	247.2	0	279.4	319.7	0	
Homicide and legal intervention (E960-E978)	219	201	212	6,738	8,865	107.6	129.6	0	130.7	158.3	0	
..Homicide (E960-E969)	216	198	209	6,660	8,698	106.8	129.3	0	129.6	157.9	0	
All injury (E800-E999)	3,915	2,520	2,881	71,345	98,223	1026.9	1078.6	0	1303.9	1347.4	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	865	660	741	20,792	27,765	330.4	435.2	1	407.8	526.4	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	441	96	160	2,062	3,332	27.9	24.8	0	40.2	35.4	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	422	360	390	10,077	13,827	149.0	215.4	1	188.9	268.8	1	
...Suicide (E955.0-.4)	288	227	256	5,733	8,145	77.6	109.6	1	101.9	140.8	1	
...Homicide (E965.0-.4)	121	120	121	4,017	5,225	66.8	88.5	0	81.1	107.3	0	
All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	782	720	749	19,034	26,387	240.9	94.4	1	312.0	124.2	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	428	308	384	4,855	8,355	52.6	56.9	0	85.5	92.6	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	677	643	656	17,442	23,924	220.8	50.1	1	283.5	65.3	1	

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



**CONNECTICUT White Resident DEATHS, 1996-1998**  
**Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)**  
**for selected causes of death by sex of decedent**  
**\*\*\* Age-adjusted using 1940 U.S. standard population \*\*\***

	Male											
	Number of Deaths, 1996-1998				Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1996-1998	1989-1991	Significance	1996-1998	1989-1991	Significance	
<b>*****Causes of Death*****</b>												
..Asthma (493)	39	26	29	565	838	15.9	11.2	0	21.6	15.9	0	
Chronic liver disease and cirrhosis (571)	509	306	424	4,412	8,057	97.5	101.7	0	168.3	178.5	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	460	59	137	835	1,715	21.6	22.5	0	38.6	42.5	0	
..Nephritis-related (580-589)	2,253	354	812	4,373	9,888	102.9	131.0	0	209.0	260.5	1	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	556	77	174	1,117	2,274	28.8	22.4	0	51.2	43.6	0	
Unintentional injuries (E800-E949)	1,789	1,181	1,348	34,146	46,753	971.1	1014.6	0	1231.6	1257.1	0	
..Motor vehicle accidents (E810-E825)	561	454	499	14,287	19,045	443.2	651.5	1	548.4	788.6	1	
..Falls and fall related injuries (E880-E888)	314	71	119	1,543	2,498	42.7	36.7	0	61.9	52.0	0	
..Drowning (E830,E832,E910)	58	52	54	1,916	2,446	58.6	73.2	0	70.0	90.6	0	
..Residential fires (E890-E899)	45	31	38	902	1,244	25.7	40.1	0	32.6	48.7	0	
..Suicide (E950-E959)	586	478	525	12,245	17,268	332.0	411.1	1	434.3	529.9	1	
Homicide and legal intervention (E960-E978)	158	147	156	5,053	6,570	162.3	191.5	0	196.8	233.8	0	
..Homicide (E960-E969)	155	144	153	4,975	6,463	160.6	191.5	0	194.6	233.8	0	
All injury (E800-E999)	2,597	1,865	2,091	53,013	72,768	1504.2	1658.6	0	1913.0	2073.2	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	546	438	485	13,757	18,365	427.1	644.0	1	528.9	778.6	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	255	78	123	1,690	2,698	46.4	41.6	0	66.6	57.7	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	370	310	339	8,787	12,030	258.3	362.8	1	327.7	453.6	1	
...Suicide (E955.0-.4)	260	200	229	5,100	7,243	138.0	190.7	1	181.7	245.7	1	
...Homicide (E965.0-.4)	98	98	98	3,387	4,367	111.8	141.2	0	135.0	170.9	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	566	529	546	14,250	19,627	360.1	142.3	1	464.4	185.7	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	311	234	280	3,730	6,310	81.9	83.6	0	131.8	135.8	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	479	466	471	12,912	17,585	326.1	71.9	1	416.5	92.7	1	

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



CONNECTICUT White Resident DEATHS, 1996-1998  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Female											
	Number of Deaths, 1996-1998				Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1996-1998	1989-1991	Significance	1996-1998	1989-1991	Significance	
<b>*****Causes of Death*****</b>												
..Asthma (493)	100	39	53	583	1,028	13.2	6.6	0	21.4	11.3	0	
Chronic liver disease and cirrhosis (571)	311	115	199	1,603	3,158	35.0	41.2	0	63.0	75.0	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	491	49	130	677	1,539	16.3	12.3	0	31.2	25.2	0	
..Nephritis-related (580-589)	1,988	253	610	3,250	7,312	77.1	79.9	0	148.4	159.4	0	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	565	59	159	834	1,869	20.3	12.7	0	38.2	26.7	0	
Unintentional injuries (E800-E949)	1,047	426	541	12,225	16,948	372.8	327.5	0	470.8	405.9	0	
..Motor vehicle accidents (E810-E825)	325	224	260	7,070	9,460	232.2	221.0	0	285.4	269.0	0	
..Falls and fall related injuries (E880-E888)	287	16	46	322	607	7.6	7.3	0	12.3	12.3	0	
..Drowning (E830,E832,E910)	12	7	9	252	332	.	15.8	.	.	18.3	.	
..Residential fires (E890-E899)	35	21	26	609	836	17.6	29.3	0	22.0	35.8	0	
Suicide (E950-E959)	173	139	156	3,523	5,005	94.1	81.5	0	123.1	108.5	0	
Homicide and legal intervention (E960-E978)	61	54	56	1,685	2,235	50.5	66.0	0	62.0	81.1	0	
..Homicide (E960-E969)	61	54	56	1,685	2,235	50.5	65.4	0	62.0	80.4	0	
All injury (E800-E999)	1,318	655	790	18,333	25,455	539.4	487.7	0	684.9	612.1	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	319	222	256	7,035	9,460	231.1	220.7	0	283.9	268.2	0	
..All falls (E880-E886,E888,E957,E968.1,E987)	186	18	37	372	635	9.1	7.7	0	13.7	13.0	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-	4	52	50	51	1,290	1,798	36.8	65.0	0	47.2	81.5	0
...Suicide (E955.0-.4)	28	27	27	633	903	16.3	27.5	0	21.5	35.5	0	
....Homicide (E965.0-.4)	23	22	23	630	858	19.9	34.3	0	25.0	42.1	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	216	191	203	4,785	6,760	120.7	46.4	1	158.6	62.9	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-	3,790.3,E860)	117	74	104	1,125	2,045	23.8	30.8	0	40.5	51.2	0
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-	5,E962.0,E980.0-.5)	198	177	185	4,530	6,340	114.6	28.3	1	149.4	38.1	1

\* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



CONNECTICUT White Resident DEATHS, 1996-1998  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Both Sexes												
	Number of Deaths, 1996-1998			Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75				
	All	< 65	< 75	< 65	< 75	1996-1998	1989-1991	Significance	1996-1998	1989-1991	Significance		
<b>*****Causes of Death*****</b>													
..Asthma (493)	139	65	82	1,148	1,865	13.9	8.9	0	20.9	13.8	0		
Chronic liver disease and cirrhosis (571)	820	421	623	6,015	11,215	62.9	66.8	0	113.7	123.6	0		
Nephritis, nephrotic syndrome/nephrosis (580-589)	951	108	267	1,511	3,254	18.7	17.4	0	36.4	35.8	0		
..Nephritis-related (580-589)	4,241	607	1,422	7,623	17,200	88.7	103.1	0	186.5	218.5	1		
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	1,121	136	333	1,950	4,143	24.3	17.6	0	46.6	37.4	0		
Unintentional injuries (E800-E949)	2,836	1,607	1,889	46,371	63,701	631.7	650.0	0	791.4	794.9	0		
..Motor vehicle accidents (E810-E825)	886	678	759	21,357	28,505	323.7	421.0	1	394.3	502.5	1		
..Falls and fall related injuries (E880-E888)	601	87	165	1,865	3,165	23.8	22.0	0	35.8	32.5	0		
..Drowning (E830,E832,E910)	70	59	63	2,168	2,778	33.3	44.7	0	39.1	53.6	0		
..Residential fires (E890-E899)	80	52	64	1,510	2,080	22.4	36.3	0	27.9	43.5	0		
Suicide (E950-E959)	759	617	681	15,768	22,273	192.4	224.7	0	250.3	289.3	0		
Homicide and legal intervention (E960-E978)	219	201	212	6,738	8,865	103.5	122.2	0	123.8	147.0	0		
..Homicide (E960-E969)	216	198	209	6,660	8,698	102.8	122.0	0	122.9	146.7	0		
All injury (E800-E999)	3,915	2,520	2,881	71,345	98,223	953.7	1020.6	0	1199.2	1261.0	0		
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	865	660	741	20,792	27,765	315.7	417.2	1	384.6	497.2	1		
..All falls (E880-E886,E888,E957,E968.1,E987)	441	96	160	2,062	3,332	26.0	24.4	0	38.2	35.3	0		
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	422	360	390	10,077	13,827	135.2	199.8	1	170.1	246.5	1		
...Suicide (E955.0-.4)	288	227	256	5,733	8,145	68.8	99.8	1	90.3	127.4	1		
...Homicide (E965.0-.4)	121	120	121	4,017	5,225	62.3	83.4	0	74.4	99.5	0		
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	782	720	749	19,034	26,387	208.2	83.6	1	267.4	109.8	1		
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	428	308	384	4,855	8,355	48.6	52.4	0	81.5	88.3	0		
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	677	643	656	17,442	23,924	189.5	42.6	1	240.7	55.4	1		

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



CONNECTICUT White Resident DEATHS, 1996-1998  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Male											
	Number of Deaths, 1996-1998			Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1996-1998	1989-1991	Significance	1996-1998	1989-1991	Significance	
<b>*****Causes of Death*****</b>												
..Asthma (493)	39	26	29	565	838	14.9	10.9	0	20.3	15.7	0	
Chronic liver disease and cirrhosis (571)	509	306	424	4,412	8,057	93.6	95.9	0	167.0	175.6	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	460	59	137	835	1,715	21.4	22.4	0	40.2	44.8	0	
..Nephritis-related (580-589)	2,253	354	812	4,373	9,888	101.9	127.6	0	220.4	272.5	1	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	556	77	174	1,117	2,274	28.4	22.4	0	53.1	46.3	0	
Unintentional injuries (E800-E949)	1,789	1,181	1,348	34,146	46,753	897.8	970.2	0	1127.7	1188.1	0	
..Motor vehicle accidents (E810-E825)	561	454	499	14,287	19,045	418.6	622.0	1	511.3	741.7	1	
..Falls and fall related injuries (E880-E888)	314	71	119	1,543	2,498	40.0	36.5	0	59.4	52.4	0	
..Drowning (E830,E832,E910)	58	52	54	1,916	2,446	57.3	72.2	0	67.3	87.9	0	
..Residential fires (E890-E899)	45	31	38	902	1,244	25.4	41.9	0	31.9	50.3	0	
Suicide (E950-E959)	586	478	525	12,245	17,268	298.0	374.0	1	388.6	480.0	1	
Homicide and legal intervention (E960-E978)	158	147	156	5,053	6,570	154.8	180.7	0	184.8	217.4	0	
..Homicide (E960-E969)	155	144	153	4,975	6,463	153.4	180.7	0	183.0	217.4	0	
All injury (E800-E999)	2,597	1,865	2,091	53,013	72,768	1384.0	1562.0	1	1744.1	1931.6	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	546	438	485	13,757	18,365	403.7	614.7	1	493.7	732.0	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	255	78	123	1,690	2,698	43.2	41.1	0	63.4	57.5	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-	4,4,E970,E985.0-.4)	370	310	339	8,787	12,030	233.6	337.7	1	294.3	417.8	1
....Suicide (E955.0-.4)	260	200	229	5,100	7,243	121.9	174.4	1	160.8	223.7	1	
....Homicide (E965.0-.4)	98	98	98	3,387	4,367	103.9	133.3	0	123.4	158.7	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	566	529	546	14,250	19,627	309.5	125.6	1	395.5	163.1	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-	3,790.3,E860)	311	234	280	3,730	6,310	75.6	76.7	0	125.2	129.1	0
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-	5,5,E962.0,E980.0-.5)	479	466	471	12,912	17,585	278.4	60.2	1	351.4	77.0	1

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



**CONNECTICUT White Resident DEATHS, 1996-1998**  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Female												
	Number of Deaths, 1996-1998				Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All		< 65	< 75	< 65		< 75	1996-	1989-	Signif-icance	1996-	1989-	Signif-icance
								1998	1991		1998	1991	
*****Causes of Death*****													
..Asthma (493)	100	39	53	583	1,028	12.6	6.8	0	21.1	11.8	0		
Chronic liver disease and cirrhosis (571)	311	115	199	1,603	3,158	33.0	38.7	0	62.5	74.4	0		
Nephritis, nephrotic syndrome/nephrosis (580-589)	491	49	130	677	1,539	16.0	12.4	0	32.7	27.2	0		
..Nephritis-related (580-589)	1,988	253	610	3,250	7,312	76.1	79.4	0	155.2	168.9	0		
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	565	59	159	834	1,869	20.2	13.0	0	40.3	29.0	0		
Unintentional injuries (E800-E949)	1,047	426	541	12,225	16,948	359.5	322.1	0	449.8	394.9	0		
..Motor vehicle accidents (E810-E825)	325	224	260	7,070	9,460	226.0	214.2	0	274.3	257.0	0		
..Falls and fall related injuries (E880-E888)	287	16	46	322	607	7.2	7.2	0	12.3	12.6	0		
..Drowning (E830,E832,E910)	12	7	9	252	332	.	16.6	.	.	18.8	.		
..Residential fires (E890-E899)	35	21	26	609	836	19.4	30.4	0	23.8	36.5	0		
Suicide (E950-E959)	173	139	156	3,523	5,005	85.2	73.4	0	110.8	97.8	0		
Homicide and legal intervention (E960-E978)	61	54	56	1,685	2,235	50.0	62.0	0	60.3	74.9	0		
..Homicide (E960-E969)	61	54	56	1,685	2,235	50.0	61.5	0	60.3	74.4	0		
All injury (E800-E999)	1,318	655	790	18,333	25,455	513.2	467.9	0	645.3	581.1	0		
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	319	222	256	7,035	9,460	225.1	213.8	0	272.9	256.2	0		
..All falls (E880-E886,E888,E957,E968.1,E987)	186	18	37	372	635	8.5	7.3	0	13.2	13.1	0		
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	52	50	51	1,290	1,798	34.1	58.8	0	43.3	72.8	0		
....Suicide (E955.0-.4)	28	27	27	633	903	14.8	24.1	0	19.4	30.8	0		
....Homicide (E965.0-.4)	23	22	23	630	858	18.8	32.1	0	23.3	38.8	0		
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	216	191	203	4,785	6,760	105.9	41.5	1	138.5	56.5	1		
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	117	74	104	1,125	2,045	22.2	28.8	0	39.2	49.3	0		
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	198	177	185	4,530	6,340	99.6	25.2	1	129.1	33.9	1		

\* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



CONNECTICUT White Resident DEATHS, 1996-1998  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Both Sexes											
	Number of Deaths, 1996-1998			Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1996-1998	1989-1991	Significance	1996-1998	1989-1991	Significance	
<b>*****Causes of Death*****</b>												
..Asthma (493)	139	65	82	1,148	1,865	15.5	9.2	0	23.1	14.5	0	
Chronic liver disease and cirrhosis (571)	820	421	623	6,015	11,215	78.2	84.0	0	134.6	146.9	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	951	108	267	1,511	3,254	20.3	18.6	0	39.2	37.9	0	
..Nephritis-related (580-589)	4,241	607	1,422	7,623	17,200	101.0	116.3	0	205.3	238.8	1	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	1,121	136	333	1,950	4,143	26.2	18.9	0	49.9	39.6	0	
Unintentional injuries (E800-E949)	2,836	1,607	1,889	46,371	63,701	639.0	611.3	0	808.1	754.9	0	
..Motor vehicle accidents (E810-E825)	886	678	759	21,357	28,505	305.6	385.7	1	375.5	463.0	1	
..Falls and fall related injuries (E880-E888)	601	87	165	1,865	3,165	25.3	21.5	0	38.1	32.3	0	
..Drowning (E830,E832,E910)	70	59	63	2,168	2,778	30.7	42.2	0	36.4	51.2	0	
..Residential fires (E890-E899)	80	52	64	1,510	2,080	21.1	32.3	0	26.7	39.3	0	
Suicide (E950-E959)	759	617	681	15,768	22,273	210.9	235.6	0	275.2	304.9	0	
Homicide and legal intervention (E960-E978)	219	201	212	6,738	8,865	96.9	117.6	0	116.8	142.5	0	
..Homicide (E960-E969)	216	198	209	6,660	8,698	96.0	117.2	0	115.6	142.0	0	
All injury (E800-E999)	3,915	2,520	2,881	71,345	98,223	978.4	992.1	0	1241.1	1237.1	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	865	660	741	20,792	27,765	297.8	382.1	1	365.9	457.9	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	441	96	160	2,062	3,332	27.8	24.1	0	41.0	35.3	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	422	360	390	10,077	13,827	139.1	197.8	1	176.0	246.1	1	
...Suicide (E955.0-.4)	288	227	256	5,733	8,145	76.1	103.9	1	100.0	133.5	1	
...Homicide (E965.0-.4)	121	120	121	4,017	5,225	58.5	78.6	0	70.3	94.4	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	782	720	749	19,034	26,387	246.8	94.3	1	317.3	123.9	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	428	308	384	4,855	8,355	62.2	66.5	0	99.6	107.0	0	
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	677	643	656	17,442	23,924	225.5	50.7	1	287.0	65.7	1	

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



**CONNECTICUT White Resident DEATHS, 1996-1998**  
**Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)**  
**for selected causes of death by sex of decedent**  
**\*\*\* Age-adjusted using 2000 U.S. standard population \*\*\***

	Male											
	Number of Deaths, 1996-1998			Years of life lost, 1996-1998		Adjusted YPLL < 65			Adjusted YPLL < 75			
	All	< 65	< 75	< 65	< 75	1996-1998	1989-1991	Significance	1996-1998	1989-1991	Significance	
<b>*****Causes of Death*****</b>												
..Asthma (493)	39	26	29	565	838	15.5	11.0	0	21.3	16.0	0	
Chronic liver disease and cirrhosis (571)	509	306	424	4,412	8,057	116.5	120.5	0	198.3	208.8	0	
Nephritis, nephrotic syndrome/nephrosis (580-589)	460	59	137	835	1,715	22.9	24.4	0	43.0	47.8	0	
..Nephritis-related (580-589)	2,253	354	812	4,373	9,888	116.9	146.4	0	243.7	300.9	1	
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	556	77	174	1,117	2,274	30.4	24.5	0	56.8	49.5	0	
Unintentional injuries (E800-E949)	1,789	1,181	1,348	34,146	46,753	926.4	920.4	0	1173.4	1137.4	0	
..Motor vehicle accidents (E810-E825)	561	454	499	14,287	19,045	400.9	571.5	1	493.8	685.2	1	
..Falls and fall related injuries (E880-E888)	314	71	119	1,543	2,498	42.0	35.7	0	62.5	52.1	0	
..Drowning (E830,E832,E910)	58	52	54	1,916	2,446	53.4	69.5	0	63.2	85.6	0	
..Residential fires (E890-E899)	45	31	38	902	1,244	24.8	36.7	0	31.7	44.6	0	
..Suicide (E950-E959)	586	478	525	12,245	17,268	326.3	389.7	0	426.8	503.1	0	
Homicide and legal intervention (E960-E978)	158	147	156	5,053	6,570	144.5	172.1	0	173.7	208.4	0	
..Homicide (E960-E969)	155	144	153	4,975	6,463	142.6	172.1	0	171.2	208.4	0	
All injury (E800-E999)	2,597	1,865	2,091	53,013	72,768	1437.6	1524.5	0	1826.0	1902.0	0	
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	546	438	485	13,757	18,365	386.3	564.7	1	476.4	675.9	1	
..All falls (E880-E886,E888,E957,E968.1,E987)	255	78	123	1,690	2,698	45.8	40.3	0	67.3	57.1	0	
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-	4,4,E970,E985.0-.4)	370	310	339	8,787	12,030	240.4	331.5	1	304.5	413.4	1
...Suicide (E955.0-.4)	260	200	229	5,100	7,243	134.7	180.0	0	177.5	232.4	0	
...Homicide (E965.0-.4)	98	98	98	3,387	4,367	97.6	124.3	0	116.4	148.8	0	
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	566	529	546	14,250	19,627	368.7	142.1	1	471.8	184.8	1	
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-	3,790.3,E860)	311	234	280	3,730	6,310	96.6	97.2	0	153.2	156.2	0
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-	5,5,E962.0,E980.0-.5)	479	466	471	12,912	17,585	333.0	73.1	1	421.2	93.3	1

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



