

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	3.5	1.32	2	1.9	1.37	5	5.1	2.29
5-9	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	0	0.0	0.00	1	1.2	1.20
20-24	6	2.9	1.17	3	2.8	1.63	3	2.9	1.68
25-29	21	8.7	1.91	14	11.6	3.10	7	5.9	2.22
30-34	59	22.3	2.90	46	34.7	5.12	13	9.8	2.72
35-39	55	22.4	3.03	48	39.4	5.68	7	5.7	2.15
40-44	38	16.6	2.69	34	30.2	5.18	4	3.4	1.71
45-49	33	17.9	3.11	28	30.9	5.83	5	5.3	2.39
50-54	22	15.3	3.27	17	24.2	5.87	5	6.8	3.05
55-59	21	16.1	3.52	17	26.9	6.53	4	6.0	2.98
60-64	24	17.6	3.60	15	23.1	5.98	9	12.6	4.20
65-69	38	29.4	4.77	22	37.9	8.08	16	22.5	5.61
70-74	55	48.7	6.56	24	49.9	10.19	31	47.7	8.57
75-79	71	81.8	9.71	34	99.5	17.07	37	70.3	11.56
80-84	59	107.5	13.99	25	134.2	26.84	34	93.8	16.08
85+	137	293.2	25.05	42	350.1	54.02	95	273.6	28.07
Total deaths	648			371			277		
Connecticut									
. Crude rate		22.1	-		26.0	-		18.3	-
. AAMR (1940 Std. mil.)		13.3	0.60		19.2	1.05		7.9	0.61
. AAMR (1970 Std. mil.)		15.4	0.63		21.1	1.12		10.3	0.68
. AAMR (2000 Std. mil.)		20.8	0.82		28.5	1.54		14.2	0.87
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		18.7	-		29.1	-		8.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.0	0.99	2	1.9	1.37	2	2.0	1.45
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	0	0.0	0.00	1	1.2	1.20
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.42	0	0.0	0.00	1	0.8	0.84
30-34	2	0.8	0.53	1	0.8	0.76	1	0.8	0.75
35-39	3	1.2	0.71	2	1.6	1.16	1	0.8	0.81
40-44	4	1.7	0.87	3	2.7	1.54	1	0.9	0.86
45-49	5	2.7	1.21	4	4.4	2.20	1	1.1	1.07
50-54	9	6.3	2.09	5	7.1	3.19	4	5.5	2.73
55-59	7	5.4	2.03	5	7.9	3.54	2	3.0	2.11
60-64	13	9.5	2.65	5	7.7	3.45	8	11.2	3.96
65-69	27	20.9	4.02	14	24.1	6.45	13	18.2	5.06
70-74	41	36.3	5.66	21	43.7	9.53	20	30.8	6.88
75-79	58	66.8	8.78	27	79.0	15.21	31	58.9	10.58
80-84	53	96.6	13.26	22	118.1	25.18	31	85.5	15.36
85+	121	259.0	23.54	38	316.7	51.38	83	239.0	26.24
Total deaths	349			149			200		
Connecticut									
. Crude rate		11.9	-		10.5	-		13.2	-
. AAMR (1940 Std. mil.)		5.0	0.32		5.8	0.51		4.4	0.41
. AAMR (1970 Std. mil.)		7.6	0.42		9.0	0.74		6.8	0.52
. AAMR (2000 Std. mil.)		11.1	0.60		13.4	1.14		9.9	0.71
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.0	-		4.5	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.86	0	0.0	0.00	3	3.1	1.77
5-9	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	5	2.4	1.07	2	1.9	1.33	3	2.9	1.68
25-29	18	7.5	1.77	13	10.8	2.98	5	4.2	1.88
30-34	52	19.6	2.72	42	31.7	4.89	10	7.5	2.39
35-39	48	19.6	2.83	42	34.5	5.32	6	4.9	1.99
40-44	33	14.4	2.51	30	26.7	4.87	3	2.6	1.48
45-49	21	11.4	2.48	18	19.8	4.68	3	3.2	1.85
50-54	12	8.4	2.41	11	15.7	4.73	1	1.4	1.36
55-59	5	3.8	1.72	4	6.3	3.17	1	1.5	1.49
60-64	5	3.7	1.64	5	7.7	3.45	0	0.0	0.00
65-69	2	1.5	1.09	2	3.4	2.44	0	0.0	0.00
70-74	4	3.5	1.77	2	4.2	2.94	2	3.1	2.18
75-79	3	3.5	2.00	1	2.9	2.93	2	3.8	2.69
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	212			172			40		
Connecticut									
. Crude rate		7.2	-		12.1	-		2.6	-
. AAMR (1940 Std. mil.)		6.5	0.45		10.7	0.82		2.5	0.41
. AAMR (1970 Std. mil.)		5.6	0.39		9.1	0.71		2.2	0.37
. AAMR (2000 Std. mil.)		6.9	0.48		11.4	0.88		2.5	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.3	-		20.1	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	2.5	1.11	4	3.9	1.94	1	1.0	1.02
5-9	5	2.7	1.23	0	0.0	0.00	5	5.6	2.52
10-14	5	3.0	1.32	4	4.6	2.31	1	1.2	1.21
15-19	4	2.3	1.17	3	3.4	1.97	1	1.2	1.20
20-24	20	9.6	2.14	12	11.3	3.26	8	7.8	2.75
25-29	16	6.7	1.67	9	7.4	2.48	7	5.9	2.22
30-34	34	12.8	2.20	16	12.1	3.02	18	13.6	3.20
35-39	48	19.6	2.83	21	17.2	3.76	27	21.9	4.22
40-44	115	50.2	4.68	50	44.4	6.28	65	55.7	6.91
45-49	192	104.1	7.51	87	95.9	10.28	105	112.0	10.93
50-54	270	188.1	11.45	134	190.9	16.49	136	185.4	15.90
55-59	397	304.7	15.29	203	321.4	22.56	194	288.9	20.74
60-64	652	478.7	18.75	353	544.6	28.99	299	418.9	24.22
65-69	947	732.3	23.80	517	890.6	39.17	430	603.4	29.10
70-74	1,132	1001.4	29.76	621	1292.0	51.85	511	786.4	34.79
75-79	1,069	1232.0	37.68	533	1560.3	67.58	536	1018.8	44.01
80-84	817	1488.5	52.08	409	2195.7	108.57	408	1125.2	55.71
85+	795	1701.6	60.35	325	2709.0	150.27	470	1353.5	62.43
Total deaths	6,523			3,301			3,222		
Connecticut									
. Crude rate		222.1	-		231.6	-		213.1	-
. AAMR (1940 Std. mil.)		123.8	1.71		147.2	2.71		108.1	2.22
. AAMR (1970 Std. mil.)		163.4	2.09		201.3	3.53		139.7	2.62
. AAMR (2000 Std. mil.)		204.2	2.55		258.4	4.65		173.0	3.11
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.4	-		164.9	-		112.5	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.48	1	0.9	0.94	0	0.0	0.00
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	2	0.8	0.53	0	0.0	0.00	2	1.5	1.07
35-39	7	2.9	1.08	3	2.5	1.42	4	3.2	1.62
40-44	8	3.5	1.23	4	3.6	1.78	4	3.4	1.71
45-49	12	6.5	1.88	6	6.6	2.70	6	6.4	2.61
50-54	20	13.9	3.12	12	17.1	4.94	8	10.9	3.86
55-59	38	29.2	4.73	23	36.4	7.59	15	22.3	5.77
60-64	62	45.5	5.78	30	46.3	8.45	32	44.8	7.92
65-69	94	72.7	7.50	49	84.4	12.06	45	63.1	9.41
70-74	164	145.1	11.33	92	191.4	19.96	72	110.8	13.06
75-79	143	164.8	13.78	73	213.7	25.01	70	133.1	15.90
80-84	111	202.2	19.20	47	252.3	36.80	64	176.5	22.06
85+	126	269.7	24.03	40	333.4	52.72	86	247.7	26.71
Total deaths	789			381			408		
Connecticut									
. Crude rate		26.9	-		26.7	-		27.0	-
. AAMR (1940 Std. mil.)		13.4	0.53		16.2	0.88		11.4	0.67
. AAMR (1970 Std. mil.)		18.8	0.69		22.9	1.18		15.9	0.84
. AAMR (2000 Std. mil.)		24.5	0.88		30.1	1.60		20.9	1.05
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.4	-		16.4	-		11.3	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colon cancer (153)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.48	1	0.9	0.94	0	0.0	0.00
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	2	0.8	0.53	0	0.0	0.00	2	1.5	1.07
35-39	5	2.0	0.91	2	1.6	1.16	3	2.4	1.41
40-44	6	2.6	1.07	4	3.6	1.78	2	1.7	1.21
45-49	12	6.5	1.88	6	6.6	2.70	6	6.4	2.61
50-54	18	12.5	2.96	10	14.2	4.51	8	10.9	3.86
55-59	34	26.1	4.48	21	33.3	7.26	13	19.4	5.37
60-64	47	34.5	5.03	25	38.6	7.71	22	30.8	6.57
65-69	77	59.5	6.79	39	67.2	10.76	38	53.3	8.65
70-74	141	124.7	10.50	79	164.4	18.49	62	95.4	12.12
75-79	122	140.6	12.73	61	178.6	22.86	61	115.9	14.85
80-84	92	167.6	17.48	37	198.6	32.66	55	151.7	20.45
85+	104	222.6	21.83	33	275.1	47.88	71	204.5	24.27
Total deaths	662			319			343		
Connecticut									
. Crude rate		22.5	-		22.4	-		22.7	-
. AAMR (1940 Std. mil.)		11.3	0.49		13.7	0.81		9.6	0.61
. AAMR (1970 Std. mil.)		15.8	0.63		19.2	1.08		13.4	0.77
. AAMR (2000 Std. mil.)		20.6	0.81		25.2	1.46		17.6	0.96
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.3	-		13.6	-		9.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.8	0.58	1	0.8	0.82	1	0.8	0.81
40-44	2	0.9	0.62	0	0.0	0.00	2	1.7	1.21
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	2	1.4	0.99	2	2.8	2.01	0	0.0	0.00
55-59	4	3.1	1.53	2	3.2	2.24	2	3.0	2.11
60-64	14	10.3	2.75	4	6.2	3.09	10	14.0	4.43
65-69	16	12.4	3.09	9	15.5	5.17	7	9.8	3.71
70-74	22	19.5	4.15	13	27.0	7.50	9	13.9	4.62
75-79	19	21.9	5.02	10	29.3	9.26	9	17.1	5.70
80-84	16	29.2	7.29	8	42.9	15.18	8	22.1	7.80
85+	19	40.7	9.33	6	50.0	20.42	13	37.4	10.38
Total deaths	116			55			61		
Connecticut									
. Crude rate		3.9	-		3.9	-		4.0	-
. AAMR (1940 Std. mil.)		2.0	0.20		2.3	0.32		1.8	0.27
. AAMR (1970 Std. mil.)		2.7	0.26		3.3	0.44		2.4	0.33
. AAMR (2000 Std. mil.)		3.6	0.33		4.4	0.61		3.1	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.5	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	1	0.4	0.41	0	0.0	0.00	1	0.8	0.81
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	7	3.8	1.43	3	3.3	1.91	4	4.3	2.13
50-54	8	5.6	1.97	5	7.1	3.19	3	4.1	2.36
55-59	18	13.8	3.26	10	15.8	5.01	8	11.9	4.21
60-64	32	23.5	4.15	16	24.7	6.17	16	22.4	5.60
65-69	43	33.3	5.07	23	39.6	8.26	20	28.1	6.28
70-74	63	55.7	7.02	32	66.6	11.77	31	47.7	8.57
75-79	65	74.9	9.29	20	58.5	13.09	45	85.5	12.75
80-84	46	83.8	12.36	22	118.1	25.18	24	66.2	13.51
85+	47	100.6	14.67	17	141.7	34.37	30	86.4	15.77
Total deaths	331			149			182		
Connecticut									
. Crude rate		11.3	-		10.5	-		12.0	-
. AAMR (1940 Std. mil.)		5.7	0.35		6.4	0.56		5.2	0.45
. AAMR (1970 Std. mil.)		8.0	0.45		9.1	0.75		7.3	0.57
. AAMR (2000 Std. mil.)		10.3	0.57		11.8	1.00		9.4	0.70
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.3	-		7.5	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	1.1	0.65	2	1.5	1.07	1	0.8	0.75
35-39	2	0.8	0.58	2	1.6	1.16	0	0.0	0.00
40-44	8	3.5	1.23	5	4.4	1.99	3	2.6	1.48
45-49	6	3.3	1.33	3	3.3	1.91	3	3.2	1.85
50-54	7	4.9	1.84	5	7.1	3.19	2	2.7	1.93
55-59	4	3.1	1.53	2	3.2	2.24	2	3.0	2.11
60-64	10	7.3	2.32	9	13.9	4.63	1	1.4	1.40
65-69	19	14.7	3.37	9	15.5	5.17	10	14.0	4.44
70-74	9	8.0	2.65	7	14.6	5.50	2	3.1	2.18
75-79	11	12.7	3.82	5	14.6	6.55	6	11.4	4.66
80-84	1	1.8	1.82	1	5.4	5.37	0	0.0	0.00
85+	5	10.7	4.79	4	33.3	16.67	1	2.9	2.88
Total deaths	85			54			31		
Connecticut									
. Crude rate		2.9	-		3.8	-		2.0	-
. AAMR (1940 Std. mil.)		2.0	0.23		2.8	0.40		1.4	0.26
. AAMR (1970 Std. mil.)		2.3	0.25		3.3	0.45		1.5	0.29
. AAMR (2000 Std. mil.)		2.7	0.30		4.0	0.56		1.8	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.7	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.8	0.59	-	-	-	2	1.7	1.19
30-34	1	0.4	0.38	-	-	-	1	0.8	0.75
35-39	2	0.8	0.58	-	-	-	2	1.6	1.15
40-44	4	1.7	0.87	-	-	-	4	3.4	1.71
45-49	5	2.7	1.21	-	-	-	5	5.3	2.39
50-54	9	6.3	2.09	-	-	-	9	12.3	4.09
55-59	3	2.3	1.33	-	-	-	3	4.5	2.58
60-64	2	1.5	1.04	-	-	-	2	2.8	1.98
65-69	3	2.3	1.34	-	-	-	3	4.2	2.43
70-74	2	1.8	1.25	-	-	-	2	3.1	2.18
75-79	3	3.5	2.00	-	-	-	3	5.7	3.29
80-84	6	10.9	4.46	-	-	-	6	16.5	6.76
85+	5	10.7	4.79	-	-	-	5	14.4	6.44
Total deaths	47			-			47		
Connecticut									
. Crude rate		-	-		-	-		3.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		2.2	0.37
. AAMR (1970 Std. mil.)		-	-		-	-		2.4	0.37
. AAMR (2000 Std. mil.)		-	-		-	-		3.0	0.45
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	0	0.0	0.00	-	-	-	0	0.0	0.00
50-54	1	0.7	0.70	-	-	-	1	1.4	1.36
55-59	4	3.1	1.53	-	-	-	4	6.0	2.98
60-64	3	2.2	1.27	-	-	-	3	4.2	2.43
65-69	10	7.7	2.45	-	-	-	10	14.0	4.44
70-74	9	8.0	2.65	-	-	-	9	13.9	4.62
75-79	4	4.6	2.30	-	-	-	4	7.6	3.80
80-84	5	9.1	4.07	-	-	-	5	13.8	6.17
85+	7	15.0	5.66	-	-	-	7	20.2	7.62
Total deaths	43			-			43		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.4	0.24
. AAMR (1970 Std. mil.)		-	-		-	-		1.8	0.30
. AAMR (2000 Std. mil.)		-	-		-	-		2.2	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.5	0.48	-	-	-	1	1.0	0.97
25-29	1	0.4	0.42	-	-	-	1	0.8	0.84
30-34	2	0.8	0.53	-	-	-	2	1.5	1.07
35-39	1	0.4	0.41	-	-	-	1	0.8	0.81
40-44	4	1.7	0.87	-	-	-	4	3.4	1.71
45-49	9	4.9	1.63	-	-	-	9	9.6	3.20
50-54	7	4.9	1.84	-	-	-	7	9.5	3.61
55-59	14	10.7	2.87	-	-	-	14	20.9	5.57
60-64	20	14.7	3.28	-	-	-	20	28.0	6.27
65-69	26	20.1	3.94	-	-	-	26	36.5	7.16
70-74	24	21.2	4.33	-	-	-	24	36.9	7.54
75-79	24	27.7	5.65	-	-	-	24	45.6	9.31
80-84	25	45.5	9.11	-	-	-	25	68.9	13.79
85+	18	38.5	9.08	-	-	-	18	51.8	12.22
Total deaths	176			-			176		
Connecticut									
. Crude rate		-	-		-	-		11.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.5	0.56
. AAMR (1970 Std. mil.)		-	-		-	-		8.0	0.64
. AAMR (2000 Std. mil.)		-	-		-	-		9.6	0.74
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	5	2.7	1.21	2	2.2	1.56	3	3.2	1.85
50-54	2	1.4	0.99	0	0.0	0.00	2	2.7	1.93
55-59	2	1.5	1.09	1	1.6	1.58	1	1.5	1.49
60-64	15	11.0	2.84	10	15.4	4.88	5	7.0	3.13
65-69	16	12.4	3.09	13	22.4	6.21	3	4.2	2.43
70-74	28	24.8	4.68	24	49.9	10.19	4	6.2	3.08
75-79	35	40.3	6.82	21	61.5	13.41	14	26.6	7.11
80-84	24	43.7	8.93	13	69.8	19.36	11	30.3	9.15
85+	29	62.1	11.53	14	116.7	31.19	15	43.2	11.15
Total deaths	156			98			58		
Connecticut									
. Crude rate		5.3	-		6.9	-		3.8	-
. AAMR (1940 Std. mil.)		2.4	0.22		3.8	0.40		1.5	0.25
. AAMR (1970 Std. mil.)		3.6	0.29		5.8	0.59		2.2	0.31
. AAMR (2000 Std. mil.)		4.8	0.39		7.9	0.83		3.0	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.7	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.70	2	1.9	1.37	0	0.0	0.00
5-9	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
10-14	1	0.6	0.59	0	0.0	0.00	1	1.2	1.21
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	5	2.4	1.07	2	1.9	1.33	3	2.9	1.68
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	5	1.9	0.84	3	2.3	1.31	2	1.5	1.07
35-39	3	1.2	0.71	3	2.5	1.42	0	0.0	0.00
40-44	4	1.7	0.87	1	0.9	0.89	3	2.6	1.48
45-49	9	4.9	1.63	7	7.7	2.92	2	2.1	1.51
50-54	9	6.3	2.09	6	8.5	3.49	3	4.1	2.36
55-59	17	13.0	3.16	12	19.0	5.49	5	7.4	3.33
60-64	19	14.0	3.20	10	15.4	4.88	9	12.6	4.20
65-69	17	13.1	3.19	11	18.9	5.71	6	8.4	3.44
70-74	22	19.5	4.15	9	18.7	6.24	13	20.0	5.55
75-79	11	12.7	3.82	5	14.6	6.55	6	11.4	4.66
80-84	8	14.6	5.15	5	26.8	12.00	3	8.3	4.78
85+	7	15.0	5.66	3	25.0	14.44	4	11.5	5.76
Total deaths	141			80			61		
Connecticut									
. Crude rate		4.8	-		5.6	-		4.0	-
. AAMR (1940 Std. mil.)		3.5	0.32		4.5	0.52		2.7	0.39
. AAMR (1970 Std. mil.)		4.0	0.35		5.1	0.58		3.2	0.43
. AAMR (2000 Std. mil.)		4.5	0.39		5.9	0.67		3.5	0.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.8	-		4.6	-		3.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.02
5-9	3	1.6	0.95	0	0.0	0.00	3	3.4	1.95
10-14	2	1.2	0.84	2	2.3	1.64	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
20-24	4	1.9	0.96	3	2.8	1.63	1	1.0	0.97
25-29	2	0.8	0.59	1	0.8	0.83	1	0.8	0.84
30-34	4	1.5	0.75	1	0.8	0.76	3	2.3	1.31
35-39	3	1.2	0.71	2	1.6	1.16	1	0.8	0.81
40-44	8	3.5	1.23	6	5.3	2.18	2	1.7	1.21
45-49	6	3.3	1.33	5	5.5	2.46	1	1.1	1.07
50-54	9	6.3	2.09	4	5.7	2.85	5	6.8	3.05
55-59	10	7.7	2.43	5	7.9	3.54	5	7.4	3.33
60-64	18	13.2	3.12	11	17.0	5.12	7	9.8	3.71
65-69	25	19.3	3.87	16	27.6	6.89	9	12.6	4.21
70-74	45	39.8	5.93	27	56.2	10.81	18	27.7	6.53
75-79	25	28.8	5.76	11	32.2	9.71	14	26.6	7.11
80-84	25	45.5	9.11	14	75.2	20.09	11	30.3	9.15
85+	32	68.5	12.11	10	83.4	26.36	22	63.4	13.51
Total deaths	223			119			104		
Connecticut									
. Crude rate		7.6	-		8.4	-		6.9	-
. AAMR (1940 Std. mil.)		4.5	0.34		5.7	0.56		3.5	0.42
. AAMR (1970 Std. mil.)		5.7	0.40		7.3	0.68		4.5	0.49
. AAMR (2000 Std. mil.)		7.0	0.48		9.1	0.87		5.6	0.56
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.9	-		6.2	-		4.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.59	1	1.2	1.16	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	5	1.9	0.84	2	1.5	1.07	3	2.3	1.31
35-39	7	2.9	1.08	4	3.3	1.64	3	2.4	1.41
40-44	5	2.2	0.98	4	3.6	1.78	1	0.9	0.86
45-49	6	3.3	1.33	5	5.5	2.46	1	1.1	1.07
50-54	11	7.7	2.31	7	10.0	3.77	4	5.5	2.73
55-59	12	9.2	2.66	8	12.7	4.48	4	6.0	2.98
60-64	35	25.7	4.34	20	30.9	6.90	15	21.0	5.43
65-69	47	36.3	5.30	30	51.7	9.43	17	23.9	5.79
70-74	69	61.0	7.35	34	70.7	12.13	35	53.9	9.10
75-79	79	91.0	10.24	37	108.3	17.81	42	79.8	12.32
80-84	79	143.9	16.19	30	161.1	29.40	49	135.1	19.31
85+	80	171.2	19.14	18	150.0	35.36	62	178.6	22.68
Total deaths	437			201			236		
Connecticut									
. Crude rate		14.9	-		14.1	-		15.6	-
. AAMR (1940 Std. mil.)		7.1	0.39		8.8	0.65		5.7	0.45
. AAMR (1970 Std. mil.)		10.1	0.50		12.1	0.86		8.5	0.59
. AAMR (2000 Std. mil.)		13.6	0.66		15.8	1.15		11.8	0.78
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.9	-		12.7	-		11.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.59	1	1.2	1.16	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.8	0.59	2	1.7	1.17	0	0.0	0.00
30-34	9	3.4	1.13	5	3.8	1.69	4	3.0	1.51
35-39	13	5.3	1.47	7	5.7	2.17	6	4.9	1.99
40-44	16	7.0	1.75	12	10.7	3.08	4	3.4	1.71
45-49	22	11.9	2.54	18	19.8	4.68	4	4.3	2.13
50-54	33	23.0	4.00	21	29.9	6.53	12	16.4	4.72
55-59	59	45.3	5.89	34	53.8	9.23	25	37.2	7.45
60-64	135	99.1	8.53	77	118.8	13.54	58	81.3	10.67
65-69	273	211.1	12.78	149	256.7	21.03	124	174.0	15.63
70-74	338	299.0	16.26	164	341.2	26.64	174	267.8	20.30
75-79	382	440.2	22.52	186	544.5	39.92	196	372.6	26.61
80-84	349	635.9	34.04	137	735.5	62.84	212	584.7	40.16
85+	459	982.4	45.86	130	1083.6	95.04	329	947.5	52.24
Total deaths	2,091			943			1,148		
Connecticut									
. Crude rate		71.2	-		66.2	-		75.9	-
. AAMR (1940 Std. mil.)		32.2	0.79		38.8	1.33		27.1	0.96
. AAMR (1970 Std. mil.)		47.4	1.07		56.5	1.85		40.9	1.28
. AAMR (2000 Std. mil.)		64.9	1.43		76.6	2.59		56.8	1.69
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									
. The U.S. AAMR (1940 Std. mil.):									
. THE U.S. HP 2000 target:									

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.59	1	1.2	1.16	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	0	0.0	0.00	1	0.8	0.75
35-39	1	0.4	0.41	1	0.8	0.82	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	1	0.7	0.73	0	0.0	0.00	1	1.4	1.40
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	2	2.3	1.63	2	5.9	4.14	0	0.0	0.00
80-84	2	3.6	2.58	2	10.7	7.59	0	0.0	0.00
85+	1	2.1	2.14	1	8.3	8.34	0	0.0	0.00
Total deaths	9			7			2		
Connecticut									
. Crude rate		0.3	-		0.5	-		0.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. AAMR (1970 Std. mil.)		-	-		-	-		-	-
. AAMR (2000 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.5	-		0.6	-		0.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.7	0.70	0	0.0	0.00	1	1.4	1.36
55-59	1	0.8	0.77	1	1.6	1.58	0	0.0	0.00
60-64	3	2.2	1.27	2	3.1	2.18	1	1.4	1.40
65-69	11	8.5	2.56	5	8.6	3.85	6	8.4	3.44
70-74	23	20.3	4.24	6	12.5	5.10	17	26.2	6.35
75-79	29	33.4	6.21	15	43.9	11.34	14	26.6	7.11
80-84	35	63.8	10.78	13	69.8	19.36	22	60.7	12.94
85+	87	186.2	19.96	24	200.1	40.84	63	181.4	22.86
Total deaths	190			66			124		
Connecticut									
. Crude rate		6.5	-		4.6	-		8.2	-
. AAMR (1940 Std. mil.)		2.1	0.17		2.1	0.27		2.0	0.22
. AAMR (1970 Std. mil.)		3.7	0.28		3.9	0.49		3.7	0.35
. AAMR (2000 Std. mil.)		6.0	0.44		6.4	0.82		5.9	0.53
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		2.2	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	14	7.0	1.86	8	7.7	2.74	6	6.1	2.50
5-9	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	4	2.3	1.17	4	4.5	2.27	0	0.0	0.00
20-24	5	2.4	1.07	2	1.9	1.33	3	2.9	1.68
25-29	16	6.7	1.67	9	7.4	2.48	7	5.9	2.22
30-34	26	9.8	1.92	18	13.6	3.20	8	6.0	2.13
35-39	50	20.4	2.89	35	28.7	4.85	15	12.2	3.14
40-44	94	41.0	4.23	75	66.6	7.70	19	16.3	3.74
45-49	125	67.8	6.06	104	114.6	11.24	21	22.4	4.89
50-54	191	133.0	9.63	139	198.0	16.80	52	70.9	9.83
55-59	297	227.9	13.23	201	318.3	22.45	96	143.0	14.59
60-64	597	438.3	17.94	414	638.8	31.39	183	256.4	18.95
65-69	938	725.3	23.68	621	1069.7	42.93	317	444.8	24.98
70-74	1,364	1206.6	32.67	768	1597.8	57.66	596	917.2	37.57
75-79	1,697	1955.7	47.48	930	2722.4	89.27	767	1457.9	52.64
80-84	1,985	3616.6	81.17	837	4493.5	155.32	1,148	3166.1	93.44
85+	3,863	8268.2	133.03	1,099	9160.6	276.33	2,764	7959.9	151.40
Total deaths	11,269			5,265			6,004		
Connecticut									
. Crude rate		383.6	-		369.4	-		397.0	-
. AAMR (1940 Std. mil.)		155.0	1.71		209.4	3.08		112.7	1.85
. AAMR (1970 Std. mil.)		243.1	2.37		318.1	4.41		188.6	2.62
. AAMR (2000 Std. mil.)		357.1	3.38		454.3	6.52		290.5	3.78
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		186.1	-		244.8	-		139.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diseases of the heart (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	11	5.5	1.65	6	5.8	2.37	5	5.1	2.29
5-9	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	2	1.2	0.83	2	2.3	1.61	0	0.0	0.00
20-24	5	2.4	1.07	2	1.9	1.33	3	2.9	1.68
25-29	12	5.0	1.44	9	7.4	2.48	3	2.5	1.45
30-34	19	7.2	1.64	13	9.8	2.72	6	4.5	1.85
35-39	39	15.9	2.55	31	25.4	4.57	8	6.5	2.30
40-44	76	33.2	3.80	61	54.2	6.94	15	12.9	3.32
45-49	107	58.0	5.61	93	102.5	10.63	14	14.9	3.99
50-54	161	112.2	8.84	122	173.8	15.74	39	53.2	8.51
55-59	252	193.4	12.18	177	280.3	21.07	75	111.7	12.90
60-64	522	383.3	16.78	371	572.4	29.72	151	211.5	17.21
65-69	795	614.8	21.80	530	913.0	39.66	265	371.9	22.84
70-74	1,101	974.0	29.35	633	1317.0	52.34	468	720.2	33.29
75-79	1,333	1536.2	42.08	736	2154.5	79.42	597	1134.8	46.44
80-84	1,555	2833.1	71.85	670	3596.9	138.96	885	2440.8	82.05
85+	2,984	6386.8	116.92	877	7310.2	246.85	2,107	6067.8	132.19
Total deaths	8,977			4,334			4,643		
Connecticut									
. Crude rate		305.6	-		304.1	-		307.0	-
. AAMR (1940 Std. mil.)		125.9	1.56		174.7	2.83		88.0	1.64
. AAMR (1970 Std. mil.)		195.4	2.14		262.4	4.01		146.7	2.32
. AAMR (2000 Std. mil.)		284.1	3.01		371.4	5.88		224.7	3.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		148.2	-		200.7	-		106.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.02
5-9	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	4	1.7	0.83	4	3.3	1.66	0	0.0	0.00
30-34	7	2.6	1.00	6	4.5	1.85	1	0.8	0.75
35-39	25	10.2	2.04	19	15.6	3.58	6	4.9	1.99
40-44	56	24.4	3.27	45	40.0	5.96	11	9.4	2.84
45-49	85	46.1	5.00	78	86.0	9.74	7	7.5	2.82
50-54	119	82.9	7.60	92	131.1	13.66	27	36.8	7.08
55-59	190	145.8	10.58	137	216.9	18.53	53	78.9	10.84
60-64	396	290.8	14.61	297	458.2	26.59	99	138.7	13.94
65-69	594	459.3	18.85	410	706.3	34.88	184	258.2	19.03
70-74	830	734.2	25.49	482	1002.8	45.68	348	535.6	28.71
75-79	984	1134.0	36.15	555	1624.7	68.96	429	815.4	39.37
80-84	1,160	2113.5	62.05	501	2689.6	120.16	659	1817.5	70.80
85+	2,217	4745.2	100.78	647	5393.0	212.02	1,570	4521.4	114.11
Total deaths	6,669			3,273			3,396		
Connecticut									
. Crude rate		227.0	-		229.7	-		224.6	-
. AAMR (1940 Std. mil.)		93.2	1.33		132.3	2.46		62.6	1.35
. AAMR (1970 Std. mil.)		145.0	1.84		198.2	3.48		105.9	1.95
. AAMR (2000 Std. mil.)		211.0	2.60		279.3	5.08		163.8	2.83
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		117.8	-		163.5	-		81.6	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.44	1	0.9	0.89	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	2	1.4	0.99	2	2.8	2.01	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	9	6.6	2.20	3	4.6	2.67	6	8.4	3.43
65-69	15	11.6	2.99	10	17.2	5.45	5	7.0	3.14
70-74	42	37.2	5.73	15	31.2	8.06	27	41.6	8.00
75-79	60	69.1	8.93	29	84.9	15.76	31	58.9	10.58
80-84	76	138.5	15.88	32	171.8	30.37	44	121.3	18.29
85+	192	411.0	29.66	49	408.4	58.35	143	411.8	34.44
Total deaths	398			142			256		
Connecticut									
. Crude rate		13.5	-		10.0	-		16.9	-
. AAMR (1940 Std. mil.)		4.2	0.24		4.7	0.41		3.8	0.29
. AAMR (1970 Std. mil.)		7.7	0.39		8.4	0.71		7.2	0.48
. AAMR (2000 Std. mil.)		12.7	0.64		13.8	1.20		12.0	0.75
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		6.4	-		5.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.53	1	0.8	0.76	1	0.8	0.75
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	1.1	0.77	1	1.1	1.10	1	1.1	1.07
50-54	2	1.4	0.99	1	1.4	1.42	1	1.4	1.36
55-59	3	2.3	1.33	2	3.2	2.24	1	1.5	1.49
60-64	1	0.7	0.73	1	1.5	1.54	0	0.0	0.00
65-69	11	8.5	2.56	8	13.8	4.87	3	4.2	2.43
70-74	11	9.7	2.93	4	8.3	4.16	7	10.8	4.07
75-79	20	23.0	5.15	12	35.1	10.14	8	15.2	5.38
80-84	27	49.2	9.47	13	69.8	19.36	14	38.6	10.32
85+	31	66.4	11.92	6	50.0	20.42	25	72.0	14.40
Total deaths	110			49			61		
Connecticut									
. Crude rate		3.7	-		3.4	-		4.0	-
. AAMR (1940 Std. mil.)		1.5	0.17		1.9	0.29		1.2	0.20
. AAMR (1970 Std. mil.)		2.4	0.23		2.9	0.42		1.9	0.27
. AAMR (2000 Std. mil.)		3.5	0.33		4.1	0.61		3.0	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.2	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.70	1	1.0	0.97	1	1.0	1.02
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	4	1.7	0.83	0	0.0	0.00	4	3.4	1.68
30-34	5	1.9	0.84	4	3.0	1.51	1	0.8	0.75
35-39	9	3.7	1.22	4	3.3	1.64	5	4.1	1.82
40-44	15	6.5	1.69	11	9.8	2.95	4	3.4	1.71
45-49	12	6.5	1.88	7	7.7	2.92	5	5.3	2.39
50-54	21	14.6	3.19	12	17.1	4.94	9	12.3	4.09
55-59	29	22.3	4.13	13	20.6	5.71	16	23.8	5.96
60-64	51	37.4	5.24	26	40.1	7.87	25	35.0	7.00
65-69	83	64.2	7.05	49	84.4	12.06	34	47.7	8.18
70-74	169	149.5	11.50	85	176.8	19.18	84	129.3	14.10
75-79	242	278.9	17.93	125	365.9	32.73	117	222.4	20.56
80-84	302	550.2	31.66	106	569.1	55.27	196	540.6	38.61
85+	649	1389.1	54.53	164	1367.0	106.75	485	1396.7	63.42
Total deaths	1,594			608			986		
Connecticut									
. Crude rate		54.3	-		42.7	-		65.2	-
. AAMR (1940 Std. mil.)		19.7	0.58		22.3	0.96		17.7	0.73
. AAMR (1970 Std. mil.)		32.8	0.85		36.3	1.48		30.2	1.03
. AAMR (2000 Std. mil.)		50.9	1.28		55.4	2.33		47.7	1.53
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		26.8	-		29.3	-		24.6	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.41	0	0.0	0.00	1	0.8	0.81
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	1.1	0.77	2	2.2	1.56	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	6	4.6	1.88	5	7.9	3.54	1	1.5	1.49
60-64	9	6.6	2.20	8	12.3	4.36	1	1.4	1.40
65-69	27	20.9	4.02	20	34.5	7.70	7	9.8	3.71
70-74	41	36.3	5.66	26	54.1	10.61	15	23.1	5.96
75-79	45	51.9	7.73	28	82.0	15.49	17	32.3	7.84
80-84	42	76.5	11.81	19	102.0	23.40	23	63.4	13.23
85+	34	72.8	12.48	11	91.7	27.65	23	66.2	13.81
Total deaths	208			120			88		
Connecticut									
. Crude rate		7.1	-		8.4	-		5.8	-
. AAMR (1940 Std. mil.)		3.1	0.24		4.8	0.46		1.8	0.23
. AAMR (1970 Std. mil.)		4.7	0.33		7.1	0.65		2.9	0.33
. AAMR (2000 Std. mil.)		6.4	0.44		9.4	0.89		4.3	0.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.7	-		5.8	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.86	2	1.9	1.37	1	1.0	1.02
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	1.4	0.83	3	2.8	1.63	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	4	1.5	0.75	4	3.0	1.51	0	0.0	0.00
35-39	4	1.6	0.82	3	2.5	1.42	1	0.8	0.81
40-44	6	2.6	1.07	5	4.4	1.99	1	0.9	0.86
45-49	9	4.9	1.63	7	7.7	2.92	2	2.1	1.51
50-54	6	4.2	1.71	3	4.3	2.47	3	4.1	2.36
55-59	14	10.7	2.87	9	14.3	4.75	5	7.4	3.33
60-64	27	19.8	3.82	17	26.2	6.36	10	14.0	4.43
65-69	24	18.6	3.79	12	20.7	5.97	12	16.8	4.86
70-74	67	59.3	7.24	43	89.5	13.64	24	36.9	7.54
75-79	142	163.6	13.73	76	222.5	25.52	66	125.5	15.44
80-84	181	329.8	24.51	77	413.4	47.11	104	286.8	28.13
85+	554	1185.8	50.38	190	1583.7	114.90	364	1048.3	54.94
Total deaths	1,044			451			593		
Connecticut									
. Crude rate		35.5	-		31.6	-		39.2	-
. AAMR (1940 Std. mil.)		11.2	0.42		15.1	0.77		8.6	0.46
. AAMR (1970 Std. mil.)		20.3	0.65		27.2	1.28		16.4	0.72
. AAMR (2000 Std. mil.)		33.8	1.05		45.1	2.20		28.0	1.16
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.3	-		17.6	-		10.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.41	0	0.0	0.00	1	0.8	0.81
40-44	4	1.7	0.87	2	1.8	1.26	2	1.7	1.21
45-49	3	1.6	0.94	3	3.3	1.91	0	0.0	0.00
50-54	10	7.0	2.20	4	5.7	2.85	6	8.2	3.34
55-59	18	13.8	3.26	11	17.4	5.25	7	10.4	3.94
60-64	44	32.3	4.87	24	37.0	7.56	20	28.0	6.27
65-69	122	94.3	8.54	59	101.6	13.23	63	88.4	11.14
70-74	178	157.5	11.80	93	193.5	20.06	85	130.8	14.19
75-79	182	209.7	15.55	87	254.7	27.30	95	180.6	18.53
80-84	167	304.3	23.54	92	493.9	51.49	75	206.8	23.88
85+	219	468.7	31.67	104	866.9	85.00	115	331.2	30.88
Total deaths	951			482			469		
Connecticut									
. Crude rate		32.4	-		33.8	-		31.0	-
. AAMR (1940 Std. mil.)		13.8	0.50		17.9	0.86		11.4	0.61
. AAMR (1970 Std. mil.)		21.1	0.70		28.8	1.31		17.0	0.83
. AAMR (2000 Std. mil.)		29.4	0.96		42.0	1.99		23.2	1.08
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		20.1	-		27.0	-		15.4	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----COPD-related (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	1.2	0.83	2	2.3	1.61	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	4	1.6	0.82	1	0.8	0.82	3	2.4	1.41
40-44	7	3.1	1.15	2	1.8	1.26	5	4.3	1.92
45-49	13	7.0	1.95	10	11.0	3.49	3	3.2	1.85
50-54	27	18.8	3.62	12	17.1	4.94	15	20.4	5.28
55-59	51	39.1	5.48	28	44.3	8.38	23	34.3	7.14
60-64	126	92.5	8.24	72	111.1	13.09	54	75.6	10.29
65-69	245	189.5	12.10	133	229.1	19.87	112	157.2	14.85
70-74	383	338.8	17.31	207	430.7	29.93	176	270.9	20.42
75-79	450	518.6	24.45	242	708.4	45.54	208	395.4	27.41
80-84	448	816.2	38.56	245	1315.3	84.03	203	559.9	39.29
85+	513	1098.0	48.48	247	2058.8	131.00	266	766.0	46.97
Total deaths	2,272			1,204			1,068		
Connecticut									
. Crude rate		77.3	-		84.5	-		70.6	-
. AAMR (1940 Std. mil.)		32.9	0.78		44.7	1.36		25.7	0.93
. AAMR (1970 Std. mil.)		50.5	1.09		71.9	2.08		38.7	1.25
. AAMR (2000 Std. mil.)		70.6	1.49		104.9	3.14		53.2	1.64
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.44	0	0.0	0.00	1	0.9	0.86
45-49	2	1.1	0.77	2	2.2	1.56	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	1	0.8	0.77	1	1.6	1.58	0	0.0	0.00
60-64	1	0.7	0.73	0	0.0	0.00	1	1.4	1.40
65-69	7	5.4	2.05	2	3.4	2.44	5	7.0	3.14
70-74	4	3.5	1.77	1	2.1	2.08	3	4.6	2.67
75-79	1	1.2	1.15	0	0.0	0.00	1	1.9	1.90
80-84	3	5.5	3.16	0	0.0	0.00	3	8.3	4.78
85+	8	17.1	6.05	0	0.0	0.00	8	23.0	8.15
Total deaths	30			8			22		
Connecticut									
. Crude rate		1.0	-		0.6	-		1.5	-
. AAMR (1940 Std. mil.)		0.6	0.12		-	-		0.5	0.14
. AAMR (1970 Std. mil.)		0.7	0.14		-	-		0.8	0.18
. AAMR (2000 Std. mil.)		0.9	0.17		-	-		1.1	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.3	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.8	0.59	2	1.7	1.17	0	0.0	0.00
30-34	7	2.6	1.00	3	2.3	1.31	4	3.0	1.51
35-39	12	4.9	1.41	8	6.6	2.32	4	3.2	1.62
40-44	24	10.5	2.14	19	16.9	3.87	5	4.3	1.92
45-49	18	9.8	2.30	13	14.3	3.97	5	5.3	2.39
50-54	23	16.0	3.34	18	25.6	6.04	5	6.8	3.05
55-59	25	19.2	3.84	17	26.9	6.53	8	11.9	4.21
60-64	32	23.5	4.15	22	33.9	7.24	10	14.0	4.43
65-69	35	27.1	4.57	19	32.7	7.51	16	22.5	5.61
70-74	48	42.5	6.13	25	52.0	10.40	23	35.4	7.38
75-79	25	28.8	5.76	16	46.8	11.71	9	17.1	5.70
80-84	20	36.4	8.15	9	48.3	16.11	11	30.3	9.15
85+	6	12.8	5.24	2	16.7	11.79	4	11.5	5.76
Total deaths	277			173			104		
Connecticut									
. Crude rate		9.4	-		12.1	-		6.9	-
. AAMR (1940 Std. mil.)		6.7	0.43		9.5	0.75		4.2	0.46
. AAMR (1970 Std. mil.)		7.6	0.47		10.7	0.82		4.9	0.50
. AAMR (2000 Std. mil.)		8.9	0.54		12.6	0.97		5.8	0.58
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.2	-		11.7	-		5.1	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.02
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.42	1	0.8	0.83	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	3	1.2	0.71	3	2.5	1.42	0	0.0	0.00
40-44	3	1.3	0.76	3	2.7	1.54	0	0.0	0.00
45-49	4	2.2	1.08	2	2.2	1.56	2	2.1	1.51
50-54	6	4.2	1.71	4	5.7	2.85	2	2.7	1.93
55-59	6	4.6	1.88	3	4.8	2.74	3	4.5	2.58
60-64	26	19.1	3.74	17	26.2	6.36	9	12.6	4.20
65-69	21	16.2	3.54	14	24.1	6.45	7	9.8	3.71
70-74	47	41.6	6.06	23	47.9	9.98	24	36.9	7.54
75-79	54	62.2	8.47	29	84.9	15.76	25	47.5	9.50
80-84	65	118.4	14.69	31	166.4	29.89	34	93.8	16.08
85+	85	181.9	19.73	36	300.1	50.01	49	141.1	20.16
Total deaths	322			166			156		
Connecticut									
. Crude rate		11.0	-		11.6	-		10.3	-
. AAMR (1940 Std. mil.)		4.7	0.30		6.4	0.53		3.5	0.35
. AAMR (1970 Std. mil.)		7.1	0.41		10.0	0.78		5.4	0.47
. AAMR (2000 Std. mil.)		10.1	0.57		14.5	1.17		7.8	0.63
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.0	-		6.3	-		4.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	16	8.0	1.99	8	7.7	2.74	8	8.2	2.89
5-9	11	6.0	1.82	7	7.5	2.83	4	4.5	2.25
10-14	7	4.1	1.56	5	5.8	2.59	2	2.4	1.71
15-19	56	32.7	4.37	41	46.6	7.27	15	18.1	4.67
20-24	71	33.9	4.03	57	53.6	7.09	14	13.6	3.64
25-29	52	21.7	3.00	38	31.4	5.10	14	11.7	3.14
30-34	51	19.2	2.69	36	27.2	4.53	15	11.3	2.92
35-39	38	15.5	2.52	28	23.0	4.34	10	8.1	2.57
40-44	34	14.8	2.54	27	24.0	4.62	7	6.0	2.27
45-49	30	16.3	2.97	20	22.0	4.93	10	10.7	3.37
50-54	18	12.5	2.96	11	15.7	4.73	7	9.5	3.61
55-59	31	23.8	4.27	22	34.8	7.43	9	13.4	4.47
60-64	26	19.1	3.74	19	29.3	6.73	7	9.8	3.71
65-69	43	33.3	5.07	31	53.4	9.59	12	16.8	4.86
70-74	44	38.9	5.87	26	54.1	10.61	18	27.7	6.53
75-79	47	54.2	7.90	24	70.3	14.34	23	43.7	9.12
80-84	59	107.5	13.99	27	145.0	27.90	32	88.3	15.60
85+	117	250.4	23.15	52	433.4	60.11	65	187.2	23.22
Total deaths	751			479			272		
Connecticut									
. Crude rate		25.6	-		33.6	-		18.0	-
. AAMR (1940 Std. mil.)		20.1	0.84		29.1	1.43		11.7	0.89
. AAMR (1970 Std. mil.)		21.6	0.85		31.5	1.49		13.1	0.91
. AAMR (2000 Std. mil.)		24.6	0.91		36.4	1.73		15.5	0.97
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		31.0	-		45.4	-		17.1	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.0	0.99	2	1.9	1.37	2	2.0	1.45
5-9	5	2.7	1.23	4	4.3	2.14	1	1.1	1.12
10-14	3	1.8	1.02	2	2.3	1.64	1	1.2	1.21
15-19	41	24.0	3.74	28	31.8	6.01	13	15.7	4.34
20-24	57	27.2	3.61	48	45.1	6.51	9	8.7	2.92
25-29	31	12.9	2.32	19	15.7	3.61	12	10.1	2.91
30-34	32	12.1	2.13	19	14.3	3.29	13	9.8	2.72
35-39	15	6.1	1.58	10	8.2	2.59	5	4.1	1.82
40-44	13	5.7	1.57	9	8.0	2.67	4	3.4	1.71
45-49	13	7.0	1.95	7	7.7	2.92	6	6.4	2.61
50-54	3	2.1	1.21	1	1.4	1.42	2	2.7	1.93
55-59	15	11.5	2.97	11	17.4	5.25	4	6.0	2.98
60-64	5	3.7	1.64	2	3.1	2.18	3	4.2	2.43
65-69	13	10.1	2.79	9	15.5	5.17	4	5.6	2.81
70-74	7	6.2	2.34	4	8.3	4.16	3	4.6	2.67
75-79	15	17.3	4.46	7	20.5	7.74	8	15.2	5.38
80-84	10	18.2	5.76	5	26.8	12.00	5	13.8	6.17
85+	8	17.1	6.05	3	25.0	14.44	5	14.4	6.44
Total deaths	290			190			100		
Connecticut									
. Crude rate		9.9	-		13.3	-		6.6	-
. AAMR (1940 Std. mil.)		10.0	0.63		13.7	1.04		6.2	0.69
. AAMR (1970 Std. mil.)		9.7	0.60		13.4	1.01		6.1	0.67
. AAMR (2000 Std. mil.)		9.6	0.57		13.1	0.97		6.3	0.64
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		17.0	-		24.2	-		10.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	1.2	0.72	2	1.7	1.17	1	0.8	0.84
30-34	2	0.8	0.53	2	1.5	1.07	0	0.0	0.00
35-39	1	0.4	0.41	1	0.8	0.82	0	0.0	0.00
40-44	2	0.9	0.62	2	1.8	1.26	0	0.0	0.00
45-49	4	2.2	1.08	3	3.3	1.91	1	1.1	1.07
50-54	1	0.7	0.70	0	0.0	0.00	1	1.4	1.36
55-59	2	1.5	1.09	2	3.2	2.24	0	0.0	0.00
60-64	7	5.1	1.94	6	9.3	3.78	1	1.4	1.40
65-69	15	11.6	2.99	10	17.2	5.45	5	7.0	3.14
70-74	12	10.6	3.06	8	16.6	5.88	4	6.2	3.08
75-79	16	18.4	4.61	7	20.5	7.74	9	17.1	5.70
80-84	31	56.5	10.14	13	69.8	19.36	18	49.6	11.70
85+	74	158.4	18.41	28	233.4	44.11	46	132.5	19.53
Total deaths	170			84			86		
Connecticut									
. Crude rate		5.8	-		5.9	-		5.7	-
. AAMR (1940 Std. mil.)		2.2	0.20		3.2	0.38		1.4	0.20
. AAMR (1970 Std. mil.)		3.5	0.28		5.0	0.55		2.5	0.29
. AAMR (2000 Std. mil.)		5.4	0.42		7.7	0.88		4.1	0.45
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.7	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.70	1	1.0	0.97	1	1.0	1.02
5-9	2	1.1	0.78	2	2.1	1.51	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	1.2	0.83	2	2.3	1.61	0	0.0	0.00
20-24	3	1.4	0.83	3	2.8	1.63	0	0.0	0.00
25-29	3	1.2	0.72	3	2.5	1.43	0	0.0	0.00
30-34	2	0.8	0.53	2	1.5	1.07	0	0.0	0.00
35-39	2	0.8	0.58	2	1.6	1.16	0	0.0	0.00
40-44	3	1.3	0.76	3	2.7	1.54	0	0.0	0.00
45-49	2	1.1	0.77	2	2.2	1.56	0	0.0	0.00
50-54	3	2.1	1.21	3	4.3	2.47	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	2	1.5	1.04	2	3.1	2.18	0	0.0	0.00
65-69	2	1.5	1.09	2	3.4	2.44	0	0.0	0.00
70-74	1	0.9	0.88	0	0.0	0.00	1	1.5	1.54
75-79	2	2.3	1.63	2	5.9	4.14	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	4	8.6	4.28	2	16.7	11.79	2	5.8	4.07
Total deaths	35			31			4		
Connecticut									
. Crude rate		1.2	-		2.2	-		0.3	-
. AAMR (1940 Std. mil.)		1.1	0.20		2.0	0.38		-	-
. AAMR (1970 Std. mil.)		1.1	0.20		2.1	0.39		-	-
. AAMR (2000 Std. mil.)		1.2	0.20		2.3	0.42		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.7	-		2.9	-		0.5	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	3.0	1.22	3	2.9	1.68	3	3.1	1.77
5-9	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	2	1.2	0.83	1	1.1	1.14	1	1.2	1.20
20-24	2	1.0	0.68	1	0.9	0.94	1	1.0	0.97
25-29	1	0.4	0.42	0	0.0	0.00	1	0.8	0.84
30-34	1	0.4	0.38	0	0.0	0.00	1	0.8	0.75
35-39	2	0.8	0.58	2	1.6	1.16	0	0.0	0.00
40-44	2	0.9	0.62	0	0.0	0.00	2	1.7	1.21
45-49	1	0.5	0.54	0	0.0	0.00	1	1.1	1.07
50-54	2	1.4	0.99	1	1.4	1.42	1	1.4	1.36
55-59	2	1.5	1.09	1	1.6	1.58	1	1.5	1.49
60-64	1	0.7	0.73	0	0.0	0.00	1	1.4	1.40
65-69	1	0.8	0.77	1	1.7	1.72	0	0.0	0.00
70-74	7	6.2	2.34	2	4.2	2.94	5	7.7	3.44
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	2	4.3	3.03	2	16.7	11.79	0	0.0	0.00
Total deaths	35			15			20		
Connecticut									
. Crude rate		1.2	-		1.1	-		1.3	-
. AAMR (1940 Std. mil.)		1.1	0.20		1.0	0.27		1.3	0.31
. AAMR (1970 Std. mil.)		1.2	0.21		1.1	0.29		1.4	0.33
. AAMR (2000 Std. mil.)		1.2	0.20		1.2	0.32		1.3	0.30
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.8	-		1.1	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	20	11.7	2.61	17	19.3	4.68	3	3.6	2.09
20-24	35	16.7	2.83	34	32.0	5.48	1	1.0	0.97
25-29	36	15.0	2.50	29	24.0	4.46	7	5.9	2.22
30-34	41	15.5	2.42	31	23.4	4.20	10	7.5	2.39
35-39	29	11.8	2.20	23	18.9	3.93	6	4.9	1.99
40-44	27	11.8	2.27	21	18.7	4.07	6	5.1	2.10
45-49	26	14.1	2.76	23	25.4	5.29	3	3.2	1.85
50-54	15	10.4	2.70	9	12.8	4.27	6	8.2	3.34
55-59	16	12.3	3.07	13	20.6	5.71	3	4.5	2.58
60-64	24	17.6	3.60	19	29.3	6.73	5	7.0	3.13
65-69	10	7.7	2.45	8	13.8	4.87	2	2.8	1.98
70-74	8	7.1	2.50	8	16.6	5.88	0	0.0	0.00
75-79	12	13.8	3.99	9	26.3	8.78	3	5.7	3.29
80-84	15	27.3	7.06	14	75.2	20.09	1	2.8	2.76
85+	6	12.8	5.24	3	25.0	14.44	3	8.6	4.99
Total deaths	322			262			60		
Connecticut									
. Crude rate		11.0	-		18.4	-		4.0	-
. AAMR (1940 Std. mil.)		10.1	0.60		17.0	1.09		3.6	0.50
. AAMR (1970 Std. mil.)		9.7	0.56		16.5	1.05		3.4	0.47
. AAMR (2000 Std. mil.)		10.5	0.59		18.1	1.14		3.8	0.50
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.3	-		18.9	-		4.4	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.02
5-9	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	13	7.6	2.11	11	12.5	3.77	2	2.4	1.70
20-24	19	9.1	2.08	15	14.1	3.64	4	3.9	1.94
25-29	16	6.7	1.67	11	9.1	2.74	5	4.2	1.88
30-34	13	4.9	1.36	9	6.8	2.27	4	3.0	1.51
35-39	10	4.1	1.29	8	6.6	2.32	2	1.6	1.15
40-44	4	1.7	0.87	4	3.6	1.78	0	0.0	0.00
45-49	3	1.6	0.94	3	3.3	1.91	0	0.0	0.00
50-54	6	4.2	1.71	4	5.7	2.85	2	2.7	1.93
55-59	2	1.5	1.09	1	1.6	1.58	1	1.5	1.49
60-64	5	3.7	1.64	5	7.7	3.45	0	0.0	0.00
65-69	1	0.8	0.77	1	1.7	1.72	0	0.0	0.00
70-74	3	2.7	1.53	3	6.2	3.60	0	0.0	0.00
75-79	1	1.2	1.15	0	0.0	0.00	1	1.9	1.90
80-84	2	3.6	2.58	2	10.7	7.59	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	102			79			23		
Connecticut									
. Crude rate		3.5	-		5.5	-		1.5	-
. AAMR (1940 Std. mil.)		3.7	0.38		5.7	0.66		1.7	0.37
. AAMR (1970 Std. mil.)		3.4	0.36		5.3	0.63		1.5	0.34
. AAMR (2000 Std. mil.)		3.4	0.34		5.3	0.61		1.5	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.8	-		17.4	-		4.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	0	0.0	0.00	1	1.0	1.02
5-9	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	13	7.6	2.11	11	12.5	3.77	2	2.4	1.70
20-24	19	9.1	2.08	15	14.1	3.64	4	3.9	1.94
25-29	16	6.7	1.67	11	9.1	2.74	5	4.2	1.88
30-34	13	4.9	1.36	9	6.8	2.27	4	3.0	1.51
35-39	10	4.1	1.29	8	6.6	2.32	2	1.6	1.15
40-44	4	1.7	0.87	4	3.6	1.78	0	0.0	0.00
45-49	3	1.6	0.94	3	3.3	1.91	0	0.0	0.00
50-54	6	4.2	1.71	4	5.7	2.85	2	2.7	1.93
55-59	2	1.5	1.09	1	1.6	1.58	1	1.5	1.49
60-64	5	3.7	1.64	5	7.7	3.45	0	0.0	0.00
65-69	1	0.8	0.77	1	1.7	1.72	0	0.0	0.00
70-74	3	2.7	1.53	3	6.2	3.60	0	0.0	0.00
75-79	1	1.2	1.15	0	0.0	0.00	1	1.9	1.90
80-84	2	3.6	2.58	2	10.7	7.59	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	102			79			23		
Connecticut									
. Crude rate		3.5	-		5.5	-		1.5	-
. AAMR (1940 Std. mil.)		3.7	0.38		5.7	0.66		1.7	0.37
. AAMR (1970 Std. mil.)		3.4	0.36		5.3	0.63		1.5	0.34
. AAMR (2000 Std. mil.)		3.4	0.34		5.3	0.61		1.5	0.32
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.7	-		17.1	-		4.5	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.0	0.99	2	1.9	1.37	2	2.0	1.45
5-9	5	2.7	1.23	4	4.3	2.14	1	1.1	1.12
10-14	3	1.8	1.02	2	2.3	1.64	1	1.2	1.21
15-19	41	24.0	3.74	28	31.8	6.01	13	15.7	4.34
20-24	56	26.8	3.57	47	44.2	6.44	9	8.7	2.92
25-29	31	12.9	2.32	19	15.7	3.61	12	10.1	2.91
30-34	32	12.1	2.13	19	14.3	3.29	13	9.8	2.72
35-39	15	6.1	1.58	10	8.2	2.59	5	4.1	1.82
40-44	13	5.7	1.57	9	8.0	2.67	4	3.4	1.71
45-49	12	6.5	1.88	6	6.6	2.70	6	6.4	2.61
50-54	2	1.4	0.99	1	1.4	1.42	1	1.4	1.36
55-59	13	10.0	2.77	9	14.3	4.75	4	6.0	2.98
60-64	5	3.7	1.64	2	3.1	2.18	3	4.2	2.43
65-69	12	9.3	2.68	8	13.8	4.87	4	5.6	2.81
70-74	7	6.2	2.34	4	8.3	4.16	3	4.6	2.67
75-79	15	17.3	4.46	7	20.5	7.74	8	15.2	5.38
80-84	10	18.2	5.76	5	26.8	12.00	5	13.8	6.17
85+	8	17.1	6.05	3	25.0	14.44	5	14.4	6.44
Total deaths	284			185			99		
Connecticut									
. Crude rate		9.7	-		13.0	-		6.5	-
. AAMR (1940 Std. mil.)		9.8	0.62		13.4	1.03		6.2	0.69
. AAMR (1970 Std. mil.)		9.5	0.60		13.0	1.00		6.0	0.66
. AAMR (2000 Std. mil.)		9.4	0.56		12.7	0.95		6.2	0.64
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.7	-		23.6	-		9.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	1.2	0.72	2	1.7	1.17	1	0.8	0.84
30-34	6	2.3	0.92	4	3.0	1.51	2	1.5	1.07
35-39	1	0.4	0.41	1	0.8	0.82	0	0.0	0.00
40-44	2	0.9	0.62	2	1.8	1.26	0	0.0	0.00
45-49	5	2.7	1.21	3	3.3	1.91	2	2.1	1.51
50-54	2	1.4	0.99	0	0.0	0.00	2	2.7	1.93
55-59	2	1.5	1.09	2	3.2	2.24	0	0.0	0.00
60-64	7	5.1	1.94	6	9.3	3.78	1	1.4	1.40
65-69	14	10.8	2.89	9	15.5	5.17	5	7.0	3.14
70-74	11	9.7	2.93	7	14.6	5.50	4	6.2	3.08
75-79	14	16.1	4.31	6	17.6	7.17	8	15.2	5.38
80-84	24	43.7	8.93	10	53.7	16.98	14	38.6	10.32
85+	54	115.6	15.73	23	191.7	39.98	31	89.3	16.03
Total deaths	146			76			70		
Connecticut									
. Crude rate		5.0	-		5.3	-		4.6	-
. AAMR (1940 Std. mil.)		2.2	0.22		3.1	0.38		1.5	0.24
. AAMR (1970 Std. mil.)		3.2	0.27		4.6	0.53		2.3	0.29
. AAMR (2000 Std. mil.)		4.6	0.39		6.8	0.82		3.5	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	22	12.9	2.74	22	25.0	5.33	0	0.0	0.00
20-24	33	15.8	2.74	30	28.2	5.15	3	2.9	1.68
25-29	34	14.2	2.43	28	23.2	4.38	6	5.0	2.06
30-34	22	8.3	1.77	18	13.6	3.20	4	3.0	1.51
35-39	19	7.8	1.78	16	13.1	3.28	3	2.4	1.41
40-44	17	7.4	1.80	14	12.4	3.32	3	2.6	1.48
45-49	11	6.0	1.80	10	11.0	3.49	1	1.1	1.07
50-54	8	5.6	1.97	5	7.1	3.19	3	4.1	2.36
55-59	10	7.7	2.43	10	15.8	5.01	0	0.0	0.00
60-64	8	5.9	2.08	8	12.3	4.36	0	0.0	0.00
65-69	8	6.2	2.19	8	13.8	4.87	0	0.0	0.00
70-74	5	4.4	1.98	5	10.4	4.65	0	0.0	0.00
75-79	7	8.1	3.05	6	17.6	7.17	1	1.9	1.90
80-84	7	12.8	4.82	7	37.6	14.20	0	0.0	0.00
85+	3	6.4	3.71	2	16.7	11.79	1	2.9	2.88
Total deaths	217			191			26		
Connecticut									
. Crude rate		7.4	-		13.4	-		1.7	-
. AAMR (1940 Std. mil.)		7.3	0.52		13.1	0.99		1.7	0.35
. AAMR (1970 Std. mil.)		6.9	0.49		12.6	0.95		1.5	0.31
. AAMR (2000 Std. mil.)		7.1	0.49		13.1	0.96		1.7	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.1	-		26.4	-		4.2	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	9	5.3	1.75	9	10.2	3.41	0	0.0	0.00
20-24	17	8.1	1.97	17	16.0	3.87	0	0.0	0.00
25-29	23	9.6	2.00	19	15.7	3.61	4	3.4	1.68
30-34	14	5.3	1.41	12	9.1	2.62	2	1.5	1.07
35-39	11	4.5	1.35	10	8.2	2.59	1	0.8	0.81
40-44	14	6.1	1.63	11	9.8	2.95	3	2.6	1.48
45-49	9	4.9	1.63	8	8.8	3.12	1	1.1	1.07
50-54	4	2.8	1.39	3	4.3	2.47	1	1.4	1.36
55-59	9	6.9	2.30	9	14.3	4.75	0	0.0	0.00
60-64	6	4.4	1.80	6	9.3	3.78	0	0.0	0.00
65-69	7	5.4	2.05	7	12.1	4.56	0	0.0	0.00
70-74	4	3.5	1.77	4	8.3	4.16	0	0.0	0.00
75-79	6	6.9	2.82	5	14.6	6.55	1	1.9	1.90
80-84	5	9.1	4.07	5	26.8	12.00	0	0.0	0.00
85+	2	4.3	3.03	2	16.7	11.79	0	0.0	0.00
Total deaths	140			127			13		
Connecticut									
. Crude rate		4.8	-		8.9	-		0.9	-
. AAMR (1940 Std. mil.)		4.5	0.40		8.3	0.76		-	-
. AAMR (1970 Std. mil.)		4.2	0.37		8.0	0.73		-	-
. AAMR (2000 Std. mil.)		4.5	0.39		8.7	0.79		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		12.0	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
10-14	2	1.2	0.84	1	1.2	1.16	1	1.2	1.21
15-19	11	6.4	1.94	11	12.5	3.77	0	0.0	0.00
20-24	15	7.2	1.85	12	11.3	3.26	3	2.9	1.68
25-29	11	4.6	1.38	9	7.4	2.48	2	1.7	1.19
30-34	7	2.6	1.00	6	4.5	1.85	1	0.8	0.75
35-39	8	3.3	1.15	6	4.9	2.01	2	1.6	1.15
40-44	2	0.9	0.62	2	1.8	1.26	0	0.0	0.00
45-49	2	1.1	0.77	2	2.2	1.56	0	0.0	0.00
50-54	4	2.8	1.39	2	2.8	2.01	2	2.7	1.93
55-59	1	0.8	0.77	1	1.6	1.58	0	0.0	0.00
60-64	2	1.5	1.04	2	3.1	2.18	0	0.0	0.00
65-69	1	0.8	0.77	1	1.7	1.72	0	0.0	0.00
70-74	1	0.9	0.88	1	2.1	2.08	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	3.6	2.58	2	10.7	7.59	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	70			59			11		
Connecticut									
. Crude rate		2.4	-		4.1	-		0.7	-
. AAMR (1940 Std. mil.)		2.6	0.33		4.4	0.60		-	-
. AAMR (1970 Std. mil.)		2.4	0.31		4.2	0.57		-	-
. AAMR (2000 Std. mil.)		2.4	0.29		4.0	0.53		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.5	-		12.6	-		2.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All poisoning (E850-E869, E950-E952, E962, E972, E980-E982)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.59	1	1.2	1.16	0	0.0	0.00
15-19	6	3.5	1.43	3	3.4	1.97	3	3.6	2.09
20-24	12	5.7	1.65	9	8.5	2.82	3	2.9	1.68
25-29	18	7.5	1.77	14	11.6	3.10	4	3.4	1.68
30-34	25	9.4	1.89	17	12.8	3.11	8	6.0	2.13
35-39	24	9.8	2.00	18	14.8	3.48	6	4.9	1.99
40-44	21	9.2	2.00	18	16.0	3.77	3	2.6	1.48
45-49	15	8.1	2.10	14	15.4	4.12	1	1.1	1.07
50-54	10	7.0	2.20	4	5.7	2.85	6	8.2	3.34
55-59	8	6.1	2.17	4	6.3	3.17	4	6.0	2.98
60-64	8	5.9	2.08	5	7.7	3.45	3	4.2	2.43
65-69	1	0.8	0.77	0	0.0	0.00	1	1.4	1.40
70-74	2	1.8	1.25	0	0.0	0.00	2	3.1	2.18
75-79	2	2.3	1.63	0	0.0	0.00	2	3.8	2.69
80-84	3	5.5	3.16	2	10.7	7.59	1	2.8	2.76
85+	3	6.4	3.71	1	8.3	8.34	2	5.8	4.07
Total deaths	159			110			49		
Connecticut									
. Crude rate		5.4	-		7.7	-		3.2	-
. AAMR (1940 Std. mil.)		5.1	0.42		7.2	0.70		3.0	0.45
. AAMR (1970 Std. mil.)		4.6	0.38		6.4	0.63		2.8	0.43
. AAMR (2000 Std. mil.)		5.3	0.42		7.5	0.72		3.1	0.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.6	-		3.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	1.2	0.72	3	2.5	1.43	0	0.0	0.00
30-34	10	3.8	1.19	7	5.3	2.00	3	2.3	1.31
35-39	10	4.1	1.29	7	5.7	2.17	3	2.4	1.41
40-44	17	7.4	1.80	15	13.3	3.44	2	1.7	1.21
45-49	13	7.0	1.95	10	11.0	3.49	3	3.2	1.85
50-54	11	7.7	2.31	6	8.5	3.49	5	6.8	3.05
55-59	14	10.7	2.87	11	17.4	5.25	3	4.5	2.58
60-64	15	11.0	2.84	10	15.4	4.88	5	7.0	3.13
65-69	12	9.3	2.68	8	13.8	4.87	4	5.6	2.81
70-74	19	16.8	3.86	11	22.9	6.90	8	12.3	4.35
75-79	7	8.1	3.05	7	20.5	7.74	0	0.0	0.00
80-84	3	5.5	3.16	1	5.4	5.37	2	5.5	3.90
85+	2	4.3	3.03	1	8.3	8.34	1	2.9	2.88
Total deaths	136			97			39		
Connecticut									
. Crude rate		4.6	-		6.8	-		2.6	-
. AAMR (1940 Std. mil.)		3.7	0.33		5.6	0.58		1.9	0.33
. AAMR (1970 Std. mil.)		3.8	0.33		5.8	0.60		2.0	0.34
. AAMR (2000 Std. mil.)		4.5	0.39		6.8	0.70		2.4	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		10.9	-		3.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1991

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
 U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
 for
 Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	4	2.3	1.17	2	2.3	1.61	2	2.4	1.70
20-24	6	2.9	1.17	4	3.8	1.88	2	1.9	1.37
25-29	17	7.1	1.72	13	10.8	2.98	4	3.4	1.68
30-34	17	6.4	1.56	13	9.8	2.72	4	3.0	1.51
35-39	20	8.2	1.83	14	11.5	3.07	6	4.9	1.99
40-44	15	6.5	1.69	13	11.6	3.20	2	1.7	1.21
45-49	8	4.3	1.53	7	7.7	2.92	1	1.1	1.07
50-54	3	2.1	1.21	0	0.0	0.00	3	4.1	2.36
55-59	7	5.4	2.03	3	4.8	2.74	4	6.0	2.98
60-64	2	1.5	1.04	0	0.0	0.00	2	2.8	1.98
65-69	1	0.8	0.77	0	0.0	0.00	1	1.4	1.40
70-74	1	0.9	0.88	0	0.0	0.00	1	1.5	1.54
75-79	1	1.2	1.15	0	0.0	0.00	1	1.9	1.90
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	2	4.3	3.03	1	8.3	8.34	1	2.9	2.88
Total deaths	104			70			34		
Connecticut									
. Crude rate		3.5	-		4.9	-		2.2	-
. AAMR (1940 Std. mil.)		3.3	0.34		4.5	0.55		2.1	0.39
. AAMR (1970 Std. mil.)		2.9	0.30		3.8	0.47		2.0	0.36
. AAMR (2000 Std. mil.)		3.4	0.34		4.6	0.56		2.2	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.9	-		5.0	-		2.7	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
 . The CT HP 2000 target: No comparison is available
 . The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
 . THE U.S. HP 2000 target: CT rate is not significantly different