

Adult Major Depressive Disorder (MDD): Suicide Risk Assessment

Children Who Have Dental Decay or Cavities

Adult Major Depressive Disorder (MDD): Suicide Risk Assessment

Falls: Screening for Future Fall Risk

Maternal Depression Screening

Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan

Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention

Use of High-Risk Medications in the Elderly

Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents

Percentage of patients aged 18 years and older with a diagnosis of major depressive disorder (MDD) with a suicide risk assessment completed during the visit in which a new diagnosis or recurrent episode was identified

Percentage of patients 65 years of age and older who were screened for future fall risk during the measurement period

Percentage of patients 13 years of age and older with a new episode of alcohol and other drug (AOD) dependence who received the following. Two rates are reported. a. Percentage of patients who initiated treatment within 14 days of the diagnosis b. Percentage of patients who initiated treatment and who had two or more additional services with an AOD diagnosis within 30 days of the initiation visit

Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous twelve months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter
Normal Parameters: Age 18 years and older BMI => 18.5 and < 25 kg/m²

Percentage of patients aged 12 years and older screened for depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen